



XXVIIth International Conference on Ultrarelativistic Nucleus-Nucleus Collisions
(Quark Matter 2018)

Hadronic resonances, strange and multi-strange particle production in Xe-Xe and Pb-Pb collisions with ALICE at the LHC

D.S.D Albuquerque¹, for the ALICE Collaboration

Campinas State University, Brazil

Abstract

We present measurements of hadronic resonance, strange and multi-strange particle production in collisions of Xe-Xe and Pb-Pb at the center-of-mass energies of $\sqrt{s_{NN}} = 5.44$ and 5.02 TeV, respectively, by the ALICE collaboration at the LHC. Particle ratios are presented as a function of multiplicity for K_s^0 , Λ , Ξ^- , Ξ^+ , Ω^- , $\bar{\Omega}^+$, $\rho(770)^0$, $K^*(892)^0$, $\phi(1020)$ and $\Lambda(1520)$. Our results are discussed and compared with predictions of QCD-inspired event generators. Additionally, comparisons with lower energy measurements and smaller systems are also presented.

Keywords: LHC, ALICE, Heavy-Ion Collisions, QGP, Strangeness, Hadronic Resonances

1. Introduction

The measurement of strange and resonance particle production in relativistic heavy-ion collisions is of great interest to investigate the properties of hadronic matter under extreme conditions. The enhanced production of strange and multi-strange hadrons with respect to non-strange particles was historically considered as one of the signatures of the formation of a partonic phase during the evolution of the system created in such collisions [1]. Moreover, short-lived hadronic resonances have a lifetime shorter or comparable to the lifetime of the fireball [2], and thus can undergo re-scattering and regeneration processes [3, 4], causing their yield to deviate from expected thermal model predictions [5, 6, 7, 8]. Therefore, the study of resonances with different lifetimes is interesting to characterize the late stages of the fireball evolution. The ALICE detector is especially suited for the study of the Quark-Gluon Plasma (QGP) via relativistic heavy-ion collisions because of its exceptional particle identification capabilities. Recently, a systematic study was performed with a wide range of strange particles and hadronic resonances being measured in different colliding systems and at various energies, including the new Xe-Xe data sample collected in late 2017. The full description of the ALICE detector can be found in [9, 10] and for further information regarding the techniques used for the measurements reported here see [11, 12, 13, 14, 15].

Email address: dsilvade@cern.ch

2. Strangeness Enhancement

The enhancement of strange particles in heavy-ion collisions was historically quantified by calculating the strange particle yield per participant pair, as estimated with the Glauber model [16], and comparing the result to the values obtained in proton–proton (pp) collisions at the same center-of-mass energy [12]. In recent results, we see that this might not be the best approach, since the production rate of strange particles in pp collisions also increases with the event multiplicity [17].




Fig. 1. Ratio to pion of integrated yields for p , K_s^0 , Λ , ϕ , Ξ and Ω . The evolution with multiplicity at mid-rapidity, $\langle dN_{ch}/d\eta \rangle_{|\eta| < 0.5}$, is reported for several systems and energies, including pp at $\sqrt{s} = 7$ TeV [17, 18], p-Pb at $\sqrt{s_{NN}} = 5.02$ TeV [19, 20, 21], and also the ALICE preliminary results for pp at $\sqrt{s} = 13$ TeV, Xe-Xe at $\sqrt{s_{NN}} = 5.44$ TeV and Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV are included for comparison. Error bars show the statistical uncertainty, whereas the empty boxes show the total systematic uncertainty.

A better way to quantify this enhancement would be to normalize the total yield of strange particles to the yield of pions. Moreover, to overcome the difference in colliding systems and energies, the yield ratios are studied as a function of charged particle multiplicity at mid-rapidity. The evolution of this ratio with multiplicity is reported in Figure 1 for p , K_s^0 , Λ , ϕ , Ξ and Ω for all the available colliding systems.

We observe a smooth evolution with multiplicity across all systems, from low multiplicity pp to central AA collisions. Furthermore, despite differences in initial conditions and collision energy, p-Pb results follow the trend observed in pp collisions. The enhancement is observed to be more pronounced for particles with a larger strangeness content, with the ϕ exhibiting a behavior that is intermediate between K_s^0 and Ξ . A saturation value seems to be reached in AA collisions for $\langle dN_{ch}/d\eta \rangle_{|\eta| < 0.5} \gtrsim 100$, in close agreement with predictions from statistical hadronization models (SHMs) [8]. Ultimately, the new measurements in Xe-Xe collisions further corroborate the observation that, for a given particle species and multiplicity, particle ratios are independent of energy and system.

3. Resonance yields and properties of the hadronic phase

The measured hadronic resonance yields may be influenced by several factors: initial yield at chemical freeze-out, resonance lifetime, scattering cross-section of resonance decay daughters and lifetime of the hadronic phase of the system evolution. Therefore, a comparison of measured resonance yield to the production rate of its stable counterpart can provide information about the late stage of the system evolution.

Hadronic resonances reported here are reconstructed via the detection of their strong-decay daughters. The lifetimes of the measured resonances vary significantly, as indicated in Table 1.

	$\rho(770)^0$	$K^*(892)^0$	$\Lambda(1520)$	$\phi(1020)$
Lifetime (fm/c)	1.3	4.2	12.6	46
Decay mode	$\pi^+\pi^-$	$K^+\pi^-$	pK^-	K^+K^-
B.R. (%)	~ 100	~ 66.6	22.5	48.9

Table 1. Lifetime values, reconstructed decay mode and branching ratio for $\rho(770)^0$, $K^*(892)^0$, $\Lambda(1520)$ and $\phi(1020)$.

The ratio of integrated yields is shown in Figure 2 for pp, Pb-Pb and Xe-Xe collisions, for $2\rho(770)^0/(\pi^+ + \pi^-)$, $(K^*(893)^0 + \bar{K}^*(893)^0)/(K^+ + K^-)$, $(\Lambda(1520) + \bar{\Lambda}(1520))/(\Lambda + \bar{\Lambda})$ and $2\phi(1020)/(K^+ + K^-)$, henceforth denoted by the shorthand notation ρ^0/π , K^{*0}/K , $\Lambda(1520)/\Lambda$ and ϕ/K . Short-lived resonances show a sizeable dependence on multiplicity. A clear suppression is observed for ρ^0 going from pp to Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV [15]. This evolution with multiplicity is well described by the EPOS3 event generator [22, 23, 24], which includes UrQMD [25, 26] for the late stage hadronic cascading. The attempt to describe the ρ^0/π ratio without the UrQMD component of the evolution fails to reproduce the data, with an overestimated and multiplicity-independent ρ^0/π ratio [27].




Fig. 2. Ratio of resonance and stable counterpart yields is reported as a function of multiplicity for the hadronic resonances, ρ^0/π [15], K^{*0}/K , $\Lambda(1520)/\Lambda$ [14] and ϕ/K from INEL > 0 pp, Pb-Pb and Xe-Xe collisions. The error bars show the statistical uncertainty, while the empty and dark-shaded boxes show the total systematic uncertainty and the uncorrelated contribution across multiplicity bins, respectively. A comparison to model is also reported by showing the ratios predicted by the EPOS3 model for Pb-Pb at $\sqrt{s_{NN}} = 2.76$ TeV, where the late hadronic cascading is simulated with UrQMD. STAR data are also shown for the $\Lambda(1520)/\Lambda$ ratio [28].

The same behavior is observed in the K^{*0}/K ratio, where the suppression can be observed in a wider multiplicity range with data from pp at $\sqrt{s} = 7$ TeV [29], Pb-Pb at $\sqrt{s_{NN}} = 5.02$ TeV and Xe-Xe at $\sqrt{s_{NN}} = 5.44$ TeV. Predictions from EPOS3 describe the evolution of particle ratios with multiplicity.

Despite having a lifetime almost ten times longer than the ρ^0 , new measurements of the $\Lambda(1520)$ production in Pb-Pb reveal that it is suppressed with respect to the Λ production [14], similarly to what is observed for other short-lived resonances. However, at present, EPOS3 predictions can only qualitatively describe the data, showing a similar decrease with multiplicity but with a higher $\Lambda(1520)$ yield. It is also worth noting that the results are in agreement with previous measurements by STAR [28].

Finally, there is small dependence of the ϕ/π ratio with multiplicity for all measured systems. This is

expected in the context of re-scattering, considering that the ϕ -meson lives longer than the expected fireball lifetime and therefore its decay daughters will not undergo re-scattering.

4. Conclusions

Latest results on strange, multi-strange and resonances in heavy-ion collisions are reported here, including new results from the Xe-Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV. Comparing the yield of strange particles to the yield of pions, an enhancement of strange particle production is observed as the event multiplicity increases. The enhancement is more pronounced for particles with higher strangeness content and, for a given multiplicity, there is no dependence with both the colliding system and energy.

The measurement of hadronic resonances is also performed for various systems (from pp to Pb-Pb, including Xe-Xe) and one can observe that, similarly to the ground state, particle ratios seem to depend solely on the final charged-particle density for all systems. A comparison of the resonance suppression with EPOS3 prediction indicates that re-scattering and regeneration processes are required to describe the measurements.

5. Acknowledgement

This work was supported by CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) grant number 141186/2015-1.

References

- [1] J. Rafelski and B. Muller *Phys. Rev. Lett.* **48** (1982) 1066. [Erratum: *Phys. Rev. Lett.* 56,2334(1986)].
- [2] C. Markert, R. Bellwied, and I. Vitev *Phys. Lett.* **B669** (2008) 92–97, arXiv:0807.1509 [nucl-th].
- [3] M. Bleicher and J. Aichelin *Phys. Lett.* **B530** (2002) 81–87, arXiv:hep-ph/0201123 [hep-ph].
- [4] M. Bleicher and H. Stoecker *J. Phys.* **G30** (2004) S111–S118, arXiv:hep-ph/0312278 [hep-ph].
- [5] S. Wheaton and J. Cleymans *Comput. Phys. Commun.* **180** (2009) 84–106, arXiv:hep-ph/0407174 [hep-ph].
- [6] A. Andronic, P. Braun-Munzinger, and J. Stachel *Phys. Lett.* **B673** (2009) 142–145, arXiv:0812.1186 [nucl-th]. [Erratum: *Phys. Lett.* B678,516(2009)].
- [7] M. Petran, J. Letessier, J. Rafelski, and G. Torrieri *Comput. Phys. Commun.* **185** (2014) 2056–2079, arXiv:1310.5108 [hep-ph].
- [8] J. Stachel, A. Andronic, P. Braun-Munzinger, and K. Redlich *J. Phys. Conf. Ser.* **509** (2014) 012019, arXiv:1311.4662 [nucl-th].
- [9] ALICE Collaboration, K. Aamodt *et al.* *JINST* **3** (2008) S08002.
- [10] ALICE Collaboration, B. B. Abelev *et al.* *Int. J. Mod. Phys.* **A29** (2014) 1430044, arXiv:1402.4476 [nucl-ex].
- [11] ALICE Collaboration, B. B. Abelev *et al.* *Phys. Rev. Lett.* **111** (2013) 222301, arXiv:1307.5530 [nucl-ex].
- [12] ALICE Collaboration, B. B. Abelev *et al.* *Phys. Lett.* **B728** (2014) 216–227, arXiv:1307.5543 [nucl-ex]. [Erratum: *Phys. Lett.* B734,409(2014)].
- [13] ALICE Collaboration, B. B. Abelev *et al.* *Phys. Rev.* **C91** (2015) 024609, arXiv:1404.0495 [nucl-ex].
- [14] ALICE Collaboration, S. Acharya *et al.* *Submitted to: Phys. Rev. Lett.* (2018), arXiv:1805.04361 [nucl-ex].
- [15] ALICE Collaboration, S. Acharya *et al.* *Submitted to: Phys. Rev. C.* (2018), arXiv:1805.04365 [nucl-ex].
- [16] ALICE Collaboration, B. Abelev *et al.* *Phys. Rev.* **C88** no. 4, (2013) 044909, arXiv:1301.4361 [nucl-ex].
- [17] ALICE Collaboration, J. Adam *et al.* *Nature Phys.* **13** (2017) 535–539, arXiv:1606.07424 [nucl-ex].
- [18] ALICE Collaboration, S. Acharya *et al.* *Submitted to: Phys. Rev.* (2018), arXiv:1807.11321 [nucl-ex].
- [19] ALICE Collaboration, B. B. Abelev *et al.* *Phys. Lett.* **B728** (2014) 25–38, arXiv:1307.6796 [nucl-ex].
- [20] ALICE Collaboration, J. Adam *et al.* *Eur. Phys. J.* **C76** no. 5, (2016) 245, arXiv:1601.07868 [nucl-ex].
- [21] ALICE Collaboration, J. Adam *et al.* *Phys. Lett.* **B758** (2016) 389–401, arXiv:1512.07227 [nucl-ex].
- [22] H. J. Drescher, M. Hladik, S. Ostapchenko, T. Pierog, and K. Werner *Phys. Rept.* **350** (2001) 93–289, arXiv:hep-ph/0007198 [hep-ph].
- [23] K. Werner, I. Karpenko, T. Pierog, M. Bleicher, and K. Mikhailov *Phys. Rev.* **C82** (2010) 044904, arXiv:1004.0805 [nucl-th].
- [24] K. Werner, B. Guiot, I. Karpenko, and T. Pierog *Phys. Rev.* **C89** no. 6, (2014) 064903, arXiv:1312.1233 [nucl-th].
- [25] S. A. Bass *et al.* *Prog. Part. Nucl. Phys.* **41** (1998) 255–369, arXiv:nucl-th/9803035 [nucl-th]. [Prog. Part. Nucl. Phys. 41,225(1998)].
- [26] M. Bleicher *et al.* *J. Phys.* **G25** (1999) 1859–1896, arXiv:hep-ph/9909407 [hep-ph].
- [27] A. G. Knospe, C. Markert, K. Werner, J. Steinheimer, and M. Bleicher *Phys. Rev.* **C93** no. 1, (2016) 014911, arXiv:1509.07895 [nucl-th].
- [28] STAR Collaboration, B. I. Abelev *et al.* *Phys. Rev. Lett.* **97** (2006) 132301, arXiv:nucl-ex/0604019 [nucl-ex].
- [29] ALICE Collaboration, B. Abelev *et al.* *Eur. Phys. J.* **C72** (2012) 2183, arXiv:1208.5717 [hep-ex].