



Characterisation of the bacterial microbiota of a landfill-contaminated confined aquifer undergoing intrinsic remediation

Daniel Abiriga^{a,*}, Andrew Jenkins^a, Kristian Alfsnes^b, Live S. Vestgarden^a, Harald Klempe^a

^a Department of Natural Sciences and Environmental Health, Faculty of Technology, Natural Sciences and Maritime Sciences, University of South-Eastern Norway, Bø, Norway


^b Division for Infection Control and Environmental Health, Norwegian Institute of Public Health, Oslo, Norway



HIGHLIGHTS

- The microbiome of a contaminated aquifer in southeast Norway was studied.
- Culture-dependent and culture-independent techniques were employed.
- Microbial composition were different between polluted and unpolluted aquifers.
- Community composition was different across the wells in the polluted aquifer.
- Functional prediction suggests the presence of organic and inorganic metabolisers.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 16 January 2021

Received in revised form 30 March 2021

Accepted 19 April 2021

Available online 26 April 2021

Editor: José Virgílio Cruz

Keywords:

Culture-dependent

Metabarcoding

Fluorescence microscopy

Landfill leachate

Microbial composition

Contaminated groundwater

ABSTRACT

Literature on microbiomes of landfill leachate-contaminated aquifers is scarce despite groundwater contaminations from landfills being common globally. In this study, a combination of microbiological techniques was applied to groundwater samples from an aquifer contaminated by a municipal landfill and undergoing intrinsic remediation. Groundwater samples were obtained from three multilevel sampling wells placed along the groundwater flow path in the contaminated aquifer, and additionally from a background well located in a nearby uncontaminated aquifer. The samples were subjected to chemical analysis, microbial culturing and characterisation, cell counting by fluorescence microscopy, and 16S rRNA metabarcoding. Good concordance was realised with the results from the different microbiological techniques. Samples from the uncontaminated aquifer had both lower cell density and lower microbial diversity compared to samples from the contaminated aquifer. Among the wells located in the contaminated aquifer, microbial diversity increased between the well closest to the landfill and the intermediate well but was lower at the most distant well. The majority of the cultured microbes represent taxa frequently recovered from contaminated environments, with 47% belonging to taxa with previously documented bioremediation potential. Multivariate redundancy analysis showed that the microbial composition was most similar in wells located closer to the landfill, although beta diversity analysis indicated a significant difference in microbial composition across the wells. Taken together with the results of cell counting, culture, and metabarcoding, these findings illustrate the effect of landfill leachates on microbial communities.

© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Groundwater is the main source of freshwater for drinking, agriculture and industry in many places globally (Mays and Scheibe, 2018;

* Corresponding author at: Department of Natural Sciences and Environmental Health, University of South-Eastern Norway, Gullbringvegen 36, NO-3800 Bø, Norway.
E-mail address: daniel.abiriga@usn.no (D. Abiriga).

O'Connor et al., 2018), but faces serious pollution challenges (Brad et al., 2013; Pous et al., 2018; Röling et al., 2001) of which leachate from landfills is one. All over the world, landfills have served as the ultimate destination for municipal solid wastes (Reinhard et al., 1984), and continue to do so (Eggen et al., 2010). In Norway, there was little recycling of wastes until the late 1990s and most of the wastes from households and industries were deposited in municipal solid waste landfills with no provision for treatment or containment of the resultant leachate. Revdalen Landfill represents one such historic site. It was active from 1974 to 1996, leading to the contamination of Revdalen Aquifer. Pollutants of environmental concern such as heavy metals and polycyclic aromatic hydrocarbons have been detected in Revdalen Aquifer (Abiriga et al., 2020).

Several strategies exist to reclaim contaminated aquifers. They are broadly categorised as artificial and natural. The former, which includes the conventional pump and treat are faster but require a major economic input for operation and maintenance (Hyldegaard et al., 2019). Natural attenuation such as *in situ* bioremediation, on the other hand, offers inexpensive, eco-friendly yet efficient remedies (Logeshwaran et al., 2018; Nunes et al., 2013). In addition, unlike pump and treat, *in situ* bioremediation does not generate secondary wastes. It is therefore the most widely preferred option in *in situ* remediation of groundwater (O'Connor et al., 2018). However, *in situ* remediation, particularly intrinsic bioremediation is slow, and the groundwater remains polluted for a long time, although recovery can be accelerated by amended bioremediation (Logeshwaran et al., 2018). Revdalen Aquifer is undergoing intrinsic remediation and the relevance of natural attenuation processes have been discussed previously (Abiriga et al., 2020; Abiriga et al., 2021).

Traditionally, groundwater bioremediation has been demonstrated empirically by measuring geochemical parameters, with little use of microbial data (Mouser et al., 2005). Over the years, however, it has become apparent that studying microbial community composition in addition to geochemical measurements offers a more complete picture of bioremediation (Lu et al., 2012; Pilloni et al., 2019; Röling et al., 2001). In order to make inferences about bioremediation and effectively manage the processes, an understanding of the microorganisms responsible is necessary (Alfreider et al., 2002; Dlugonski, 2016; Köchling et al., 2015; Röling et al., 2000). A number of studies have been conducted on the microbiology of contaminated aquifers (Anantharaman et al., 2016; Cozzarelli et al., 2000; Harvey et al., 1984; Hug et al., 2015; Kleikemper et al., 2005; Ludvigsen et al., 1999; Röling et al., 2000; Tischer et al., 2012; Watanabe et al., 2002). Nonetheless, this area still requires more elucidation (He et al., 2018; Meckenstock et al., 2015). Bioremediation of hydrocarbon-polluted aquifers is well documented in the literature (Dojka et al., 1998; Harvey et al., 1984; Kleikemper et al., 2005; Pickup et al., 2001; Rooney-Varga et al., 1999; Tischer et al., 2012; Watanabe et al., 2002), but there is a dearth of studies on landfill leachate contaminations. The bias may reflect high-profile cases of hydrocarbon pollutions and the potential health hazard presented by the concomitant xenobiotics, which are often toxic, mutagenic and carcinogenic (Logeshwaran et al., 2018). Moreover, hydrocarbons, at least, are easily degraded in the environment and their compositions are less complex than effluents emanating from landfills. The complicated attenuation processes in landfill-leachate-impacted groundwater makes assessments of bioremediation processes more difficult (Christensen et al., 2000) and less attractive.

Studies on the microbiology of landfill-impacted aquifers have been reported (Albrechtsen et al., 1995; Lin et al., 2007; Ludvigsen et al., 1999; Mouser et al., 2005; Röling et al., 2000). These studies have provided insights into the microbial profiles of leachate-impacted aquifers, but they employed low throughput methods that cannot capture the full microbial diversity. Fewer studies have applied high throughput sequence-based methods (Chen et al., 2017; Lu et al., 2012; Taş et al., 2018) and they have not specifically focused on microbial diversity. Thus, there is insufficient literature on the microbial diversity of landfill-perturbed aquifers. The present study aimed to delineate the microbial diversity and composition of a landfill-leachate-contaminated

confined aquifer using a combination of three microbiological techniques: culture and characterisation, cell counting by fluorescence microscopy, and Illumina MiSeq 16S rRNA metabarcoding. To the best of our knowledge, no such multi-methodological study of landfill-perturbed subsurface microbiology has been reported.

2. Materials and methods

2.1. Study area and groundwater sampling wells

Revdalen Aquifer is a glaciofluvial deposit located in Vestfold and Telemark County in southeast Norway at coordinate 59°25'58.26"N and 9°06'1.53"E (Fig. 1A). In 1958, a kettle hole was adopted as a waste disposal site and through 1958–1974, the hole was directly filled up with waste. Four successive landfill cells were later opened in the area and were filled up between 1974 and 1996. Due to the leachate leaking from the landfill, the aquifer, which was serving as a water source for the surrounding communities, became contaminated and water extraction was halted. Since the closure of the landfill in February 1997, the aquifer has been allowed to undergo monitored natural attenuation. Three multilevel monitoring wells (Fig. 1B) were established along the groundwater flow direction to monitor the groundwater quality: R1, with five levels (R101–105); R2, with four levels (R201–204) and R4, with three levels (R401–403). The increasing number in the multilevel (#101–105) depicts an increasing depth in the respective wells. Details on the multilevel systems are described elsewhere (Abiriga et al., 2020). The wells are hereafter referred to as proximal, intermediate and distal wells, respectively. In addition, a background well (R0) was established in a nearby aquifer for benchmarking the water quality. Additional information on the study site can be found elsewhere (Abiriga et al., 2020, 2021; Klempe, 2004, 2015).

2.2. Groundwater sampling and chemical analyses

Groundwater sampling procedure for proximal-distal wells has been described previously (Abiriga et al., 2020) while samples from the background well were taken using a submersible pump after purging the well volume. Sampling was conducted as per ISO 5667-11 (2009). Samples for this study were collected in spring and autumn 2018 (water chemistry and 16S rRNA metabarcoding), and spring and autumn 2018–2019 (plate count and microscopic count); with one sampling campaign per season. For dissolved oxygen analysis, samples were collected in Winkler bottles, fixed onsite and protected from direct light. For other chemical analyses, samples were collected in 500 ml polyethylene bottles. pH and electrical conductivity were determined in the field using the pH meter pH-110 (VWR International) and conductivity meter Elite CTS Tester (Thermo Scientific, Singapore), respectively. The samples were maintained at ≤4 °C using icepacks and a cooler box and transported to the laboratory at University of South-Eastern Norway. Samples for iron and manganese were filtered through 0.45 µm and acidified with nitric acid to pH ~2, while samples for total nitrogen were preserved by acidifying using sulphuric acid. All samples were stored at 4 °C until analysis. Norwegian Standards were followed for determination of dissolved oxygen (NS-ISO 5813), alkalinity (NS-EN ISO 9963-2), iron (NS 4773), and manganese (NS 4773). Major ions (ammonium, sodium, potassium, calcium, magnesium, chloride, nitrate and sulphate) were determined using Ion Chromatography DIONX ICS-1100 (Thermo Scientific, USA). Total nitrogen and total organic carbon (TOC) were determined using FIAlyzer-1000 (FIALab, USA) and TOC Fusion (Teledyne Tekmar, USA), respectively.

2.3. Microbial analyses

2.3.1. Sample handling

Groundwater samples were collected in sterile 350 ml PETE bottles (VWR, UK) without headspace and transported as described above.




Fig. 1. Location of the study site on a map of Norway (A) and the location of the landfill and sampling wells R0, R1, R2 and R4 (B) [Map source: Norwegian Geographical Survey, www.norgeskart.no, with permission]. R0 is the background well located in an uncontaminated aquifer, while wells R1, R2 and R4 are located in the contaminated aquifer placed at the proximal, intermediate and distal positions, respectively. Arrows indicate the groundwater flow direction. Green shading indicates woodland, yellow indicates farmland, and grey indicates industrial land, including the landfill and an adjacent active quarry/gravel pit. Contour interval 5 m.

Upon arrival at the microbiology laboratory, 4.5 ml portions were fixed with 2.5% (final concentration) phosphate-buffered glutaraldehyde for microscopy. For 16S rRNA metabarcoding, 300 ml of water was filtered through 0.2 μm poresize 47 mm diameter polycarbonate membrane filters. The filters were stored at $-70\text{ }^{\circ}\text{C}$ prior to DNA extraction. The remaining water was used for culturing heterotrophic bacteria. All the different sample handlings were conducted within 48 h.

2.3.2. Direct microscopic count

1.9 ml of fixed groundwater samples were stained with 5 $\mu\text{g}/\text{ml}$ 4',6-diamidino-2-phenylindole (DAPI) (Kepner and Pratt, 1994). The stained cells were filtered onto 0.2 μm black polycarbonate Nuclepore membrane filters (Sigma-Aldrich, Germany), transferred onto microscope slides and overlaid with antifade mountant oil (Citifluor AF87, EMS, PA, USA). Cells were enumerated under $\times 100$ oil objective using Olympus IX70 fluorescence microscope (Tokyo, Japan). Ten fields were counted and the average count was used to estimate bacterial density using the formula: $\text{Bacteria (Cells/ml)} = (N \times A_t) / (V_f \times A_g \times d)$, where N = average number of cells, A_t = effective area of the filter paper, V_f = volume of water sample filtered, A_g = area of the counting grid, and d = dilution factor (Kepner and Pratt, 1994). No observable cells were found in our blanks and therefore correcting for background noise due to contamination was not necessary.

2.3.3. Culturing and sequencing of heterotrophic bacteria

Preliminary assessment (data not shown) indicated dilutions $\geq 10^2$ could generate countable colonies and best growth occurred on half-strength tryptic soy agar. Thus, serial dilutions of 10^2 , 10^3 , and 10^4 were prepared for each water sample and triplicates of 100 μl aliquots were spread on half-strength tryptic soy agar (20 g/l TSA and 7 g/l agar). Plates were incubated at $15\text{ }^{\circ}\text{C}$ for ≥ 5 days. Aerobes were counted following incubation under aerobic condition. Anaerobes were enumerated in a parallel setup following incubation under anaerobic condition using GasPak with EZ Anaerobe Container System (BD, USA). Plates

from two dilutions of each sample were counted and the average was reported as colony forming units (cfu) per ml.

Based on observable colony morphologies such as shape, elevation, margin, size, and colour, colonies covering the full diversity of colony morphologies were picked and purified by repeated streaking and incubation until pure cultures were obtained. These cultures were then observed by wet field microscopy at $\times 1000$ (Olympus, CX22LED RFS1, China) to observe motility and cell shape, followed by Gram staining and determination of oxidase and catalase activity (Csuros et al., 1999). Strains were stored at $-70\text{ }^{\circ}\text{C}$ in nutrient broth (Sigma, Switzerland) supplemented with 25% glycerol. The laboratory procedure for DNA extraction and sequencing of the V3–V5 16S rRNA gene region of the pure isolates is provided in the supplementary information (Method S1).

2.3.4. 16S rRNA metabarcoding

The frozen filters were retrieved and cut into two halves using sterile surgical blades and DNA was extracted from one half using DNeasy PowerSoil Kit (Qiagen, Germany) following the manufacturer's instructions. The DNA quantity and quality were checked using Qubit® Fluorometer 3.0 (Life Technologies, Malaysia) and 2% agarose gel electrophoresis, respectively. PCR and 16S rRNA gene library preparation (Fadrosh et al., 2014) for the samples were conducted at Norwegian Sequencing Centre (<https://www.sequencing.uio.no>). Both forward and reverse oligos included Illumina-specific adaptor sequence, a 12-nucleotide barcode sequence, a heterogeneity spacer and respectively the primers 319F (5'-ACTCCTACGGGAGGCAGCAG-3') and 805R (5'-GGACTACNVGGGTWCTAAT-3') for the V3–V4 hypervariable 16S rRNA gene region. The pooled libraries were sequenced using Illumina MiSeq (600 cycles), applying the 300 bp paired-end protocol with 10% PhiX.

Sequence demultiplexing was conducted using a demultiplexer available at https://github.com/nsc-norway/triple_index-demultiplexing/tree/master/src. Barcodes and heterogeneity spacers were removed in

the process with no mismatches allowed in the barcode. Denoising (primer trimming, and removal of shorter sequences and chimeras), dereplication, merging, and clustering of sequences into amplicon sequencing variants were performed using DADA2 algorithm (Callahan et al., 2016) plug-in for QIIME2 version 2019.1.0 (Bolyen et al., 2019). Default parameters were implemented, apart from primer length (adjusted to 20 bp) and minimum sequence length of reads (adjusted to 280 bp). Taxonomic assignment was conducted using Naïve Bayes classifier algorithm trained on data from SILVA v.138 using QIIME2 version 2020.2.0.

2.3.5. Quality control

Quality control samples included in the sequencing run consisted of an elution buffer, extraction blanks and mock communities, which are described in more detail in the supplementary information (Method S2).

2.4. Data availability statement

The 87 16S rRNA gene sequences of the cultured isolates have been deposited in GenBank under Accession Nos. MT348616–MT348702. The raw sequence reads from the metabarcoding have been deposited in Sequence Read Archive under BioProject PRJNA677875 (BioSamples used in this study: SAMN16775936–SAMN16775961).

2.5. Data analysis and statistics

All data analyses were conducted in R version 4.0.2 (R Core Team, 2020). To compare water chemistry between the background sample and the contaminated water samples, one-tailed Wilcoxon rank test was used. Comparison of water quality across the wells in the contaminated aquifer was performed using Kruskal–Wallis rank sum test. These tests were chosen as most of the variables showed non-normal distribution. One-tailed paired *t*-test was used for both within-sample and overall comparison between plate and microscopic counts. Similarly, comparison between aerobic and anaerobic counts was also done using one-tailed paired *t*-test. One-way ANOVA was conducted to test for differences in aerobic and microscopic counts across the wells and a post-hoc Tukey's Honest Significant Difference for the pairwise comparisons. All the parametric tests were performed on log-transformed data. Normality of data distribution was assessed graphically using histograms and boxplots, and by Shapiro–Wilk normality test. Statistical significance was inferred at $\alpha = 0.05$.

Multivariate analyses were performed using package *vegan* in R (Oksanen et al., 2019). Prior to redundancy analysis (RDA), the OTU abundance data was pre-transformed using fourth-root transformation to reduce asymmetry of the data distribution before standardising it using Hellinger standardisation (Legendre and Gallagher, 2001). Also, because groundwater physicochemical variables are dimensionally heterogeneous, the chemistry data was standardised (centred and scaled by standard deviation) prior to RDA ordination. The difference in the community profiles of the sample groups in the RDA triplot was tested using permutational multivariate analysis of variance (PERMANOVA) on Euclidean distance. The homogeneity of group dispersion was assessed beforehand using the function *betadisper* (Anderson, 2006). For the RDA, the full dataset was used without any merging ($N = 26$: R0 = 2, R1 = 10, R2 = 8 and R4 = 6). Principal component analysis (PCA) was applied on square-root transformed and Hellinger-standardised culture-based microbial data and the physicochemical data was fitted onto the PCA using the command *envfit* in package *vegan*. Taxonomic-to-phenotypic mapping was implemented using METAGENassist (Arndt et al., 2012) and to reduce the number of categories, abundances were merged by summing across the well levels and presented as the wells instead.

3. Results

3.1. Groundwater chemistry

Most of the solutes in the groundwater were present at levels significantly above that of the background well (Table 1). Kruskal–Wallis test showed significant differences across the wells in the contaminated aquifer for eight of the fifteen variables measured (Table S1). While most of the variables showed a decrease in concentration along the groundwater flow path, a few *i.e.*, dissolved oxygen, magnesium, manganese and iron showed an increase in concentration downgradient.

3.2. Groundwater microbiology

3.2.1. Viable plate count

The cell density estimate from the plate count was in the range $1 \times 10^2 - 3.2 \times 10^5$ cfu/ml (aerobic) and $0-2.4 \times 10^5$ cfu/ml (anaerobic). Despite the comparable maximum counts from the two growth conditions, the aerobic plate count was significantly higher than the anaerobic plate counts ($t = 3.63$, $df = 37$, $p = 0.0004$). The same strains dominated under both aerobic and anaerobic conditions and isolates recovered under anaerobic condition were all facultative anaerobes. Nonetheless, some strains were isolated only under anaerobic condition. Wells in the contamination plume had higher plate counts than the background well, and the distal well had lower plate counts compared to the intermediate and proximal wells (Fig. 2). The count was significantly different across the wells ($F = 3.09$, $df = 3$, $p = 0.0357$), but pairwise comparisons between the wells did not give significant results.

3.2.2. Direct fluorescence microscopy

Microscopic counts were in the range $7 \times 10^3 - 3.5 \times 10^5$ cells/ml. Cell morphologies observed under fluorescence microscopy included small rods, long large rods, coccobacilli and vibrios. Small rods were most frequently encountered. Water samples from the proximal and intermediate wells were dominated by short rods, vibrios and elongated narrow rods, some occurring in chains. The distal well on the other hand, did not exhibit predominance of any specific cell types, an observation

Table 1

Characteristics of the groundwater chemistry measured in spring and autumn 2018. Values from the background well (R0) were used as a benchmark against which those in the proximal (R1), intermediate (R2) and distal (R4) wells were compared. All units are in mg/l, except for pH (pH units), conductivity ($\mu\text{S}/\text{cm}$) and alkalinity (mM).

	Mean	Range (min - max)		
	R0 (N = 2)	R1 (N = 10)	R2 (N = 8)	R4 (N = 6)
pH	5.1	6.3–7.7	6.6–7.1	5.9–6.2
Conductivity	33	196–251	190–220	107–190
Dissolved oxygen ^a	4.5	1.3–2.9	1.3–4.8	1.1–9.1 ^b
Sodium	1.7	4.5–5.9	5.4–5.6	4.0–5.5
Potassium	0.32	5.0–7.5	6.7–7.5	6.4–7.5
Calcium	2.1	25–34	23–26	10.7–20
Magnesium	0.48	2.4–2.8	2.9–3.2	2.7–3.6
Ammonium ^c	0.0	0.13–1.9	0.32–0.57	0.0–0.41
Iron ^d	0.02	0.02	0.02	0.02–0.09
Manganese	0.04	0.05–0.45	0.17–0.31	0.01–1.4
Alkalinity	0.06	1.2–1.9	1.2–1.7	0.77–1.3
Sulphate	2.9	8.4–25	7.4–9.7	5.7–10
Nitrate (as Nitrogen)	1.1	1.6–4.8	0.48–3.7 ^b	1.3–4.0
Chloride	1.8	3.4–4.7	5.6–6.9	3.6–4.1
Total nitrogen	0.97	2.9–4.5	3.4–3.5 ^b	2.9–3.4 ^b
Total organic carbon	4.1	2.7–5.0 ^b	2.9–5.9 ^b	1.8–6.3 ^b

Values significantly higher than the background value ($p < 0.05$) are indicated in bold face. N = number of samples.

^a Dissolved oxygen was lower than the background value, but only significantly so in R1 and R2.

^b Non-significant difference from the background value ($p > 0.05$).

^c Values below the limit of detection were treated as zero.

^d Values below the limit of detection were reported as half the limit of detection (0.02).




Fig. 2. Bacterial cell density in the groundwater samples estimated from direct microscopic count (MC) and aerobic plate count (PC); error bars are mean + standard error ($n = 4$). R0 is the background well located in an aquifer upstream of the landfill. R101–R105, R201–R204 and R401–R403 are the multilevels in R1, R2 and R4 respectively, which are located in the contaminated aquifer placed along the groundwater flow direction at the proximal, intermediate and distal positions, respectively. Asterisks indicate a significant difference.

that agrees with the more diverse colony morphology (higher richness) observed in samples from this well (see below). The total cell count was not significantly different across the four wells ($F = 1.44$, $df = 3$, $p = 0.243$), nor were the pairwise comparisons.

The distal well, which showed very low plate counts, gave a relatively higher microscopic counts. Within-sample comparisons indicate that microscopic counts were, in all but one sample, higher than plate counts, but only significantly so in four of the thirteen samples: R0, R102, R105 and R204 (Fig. 2 and Table S2). Overall, the microscopic count was higher than the plate count ($t = 6.94$, $df = 51$, $p < 0.05$) (Fig. S1).

3.2.3. Identification of isolates

Small subunit rRNA gene sequences of the pure isolates revealed higher species richness in the wells located in the contamination

plume than in the background well (Fig. S2). The distal well had the highest species richness. Eighty-seven taxa belonging to forty-six genera were identified (Fig. S2, Table S3), *Pseudomonas* being the most frequently isolated taxon (Fig. S4). Of the eighty-seven taxa, seven were found only in the background aquifer, seventy-seven found only in the contaminated aquifer and three were found in both. These three were *Rhodoferrax ferrireducens*, *Pseudomonas* sp. (2) and *Rugamonas* sp. detected in both the background and distal wells (Fig. 3, Table S3). As shown in Fig. 3, twenty-eight taxa were unique to the distal well. Not a single species was found in all the four wells, although two genera (*Pseudomonas* and *Janthinobacterium*) were common. Among the wells located in the contamination plume, only two species: *Pseudomonas veronii* and *Rhodococcus degradans* were found in all of them. The proximal-intermediate was the most similar pair of wells, with ten species in common. PCA ordination further depicts this similarity (Fig. S5).




Fig. 3. Venn diagram showing relationships among the groundwater sampling wells at genus-level classification (A) and species-level classification (B). R0 is the background well located in an aquifer upstream of the landfill. R1, R2 and R4 are wells located in the contaminated aquifer placed along the groundwater flow direction at the proximal, intermediate and distal positions, respectively.

3.2.4. 16S rRNA metabarcoding

A total of 1496 OTUs were identified by metabarcoding. 98.73% were bacteria; the rest were archaea (1.2%) or unclassified (0.07%). These were classified to 57 microbial phyla, with *Proteobacteria* (31%), *Patescibacteria* (9.2%), *Bacteroidetes* (7.8%), *Actinobacteria* (6.5%), *Chloroflexi* (6.1%), *Acidobacteria* (5.5%), *Verrucomicrobia* (5.4%) and *Firmicutes* (3.5%) as the top eight phyla in the entire OTU table.

For the sake of simplicity and in order to target taxa likely to be stable members of the microbial community, only OTUs comprising at least 2% in at least one sample are considered here. This resulted in only 68 genera being represented (Fig. 4). Of these, 15 had unresolved taxonomic classifications and 17 were OTUs of uncultured microbes. Twelve of the thirty-six (33%) phylogenetically well-resolved genera were among those isolated through the culture-based approach. These were *Brevundimonas*, *Flavobacterium*, *Janthinobacterium*, *Pedobacter*, *Phyllobacterium*, *Polaromonas*, *Pseudomonas*, *Rhodanobacter*, *Rhodoferax*, *Sphingomonas*,

Stenotrophomonas and *Undibacterium*. These genera also showed greater frequency in the PCA ordination for the cultured taxa (Fig. S5). The 12 genera accounted for 38 of the total cultures isolated (Fig. S4).

Shannon diversity measure was applied to the OTU data to infer species richness and diversity. Microbial diversity was greatest in the intermediate well, followed by the proximal well and was lowest in the distal well (Fig. 5B and Table S4). Similarly, the evenness estimate followed the same trend, indicating that the distal well was dominated by few taxa, while the intermediate well had fairly even representation of microbial taxa present at the site. However, both the highest (692) and lowest (136) OTU richness was recorded at the different levels in the intermediate well. This resulted in a larger spread of OTU richness in the intermediate well (Fig. 5A). A count of species (at genus level) (Fig. 5C) indicates that a core of 137 genera were common to all the sampling wells, while a further 153 genera were common to wells of the contaminated aquifer. The number of unique genera varied between




Fig. 4. OTUs greater than 2% of the total reads in at least one of the samples. R0 is the background well located in an aquifer upstream of the landfill. R101–R105, R201–R204 and R401–R403 are the multilevels in R1, R2 and R4 respectively, which are located in the contaminated aquifer placed along the groundwater flow direction at the proximal, intermediate and distal positions, respectively. For clarity, the relative abundances for spring and autumn were merged by summing so the total per sample would be >100% but <200% since only OTUs greater than 2% are shown.




Fig. 5. OTU richness **(A)**, Shannon diversity index **(B)** (error bars are \pm SD) and species count at genus level for the full dataset **(C)** recorded in the sampling wells R0, R1, R2 and R4. For figs. **A–C**, $N = 2$ (R0), 10 (R1), 8 (R2) and 6 (R4) (Table S4). R0 is the background well located in an aquifer upstream of the landfill. R1, R2 and R4 are the wells located in the contaminated aquifer placed along the groundwater flow direction at the proximal, intermediate and distal positions, respectively.

24 in the background well and 90 in the distal well. The number of genera shared between just two wells varied from 4 (background/proximal) to 73 (distal/intermediate).

Multivariate analysis using RDA showed that microbial composition varied spatially, with the intermediate and proximal wells clustering next to each other as was observed with the culture-based method (Fig. 6 and Fig. S5). PERMANOVA showed that the microbial composition across the wells was significantly different ($F = 4.58$, $df = 3$, $p = 0.001$). Three canonical axes were significant ($p = 0.001$, 999

permutations) and the model explained 34.9% (adjusted R^2) of the variation in microbial composition. The proximal and the intermediate wells correlated positively with RDA1, while the distal and background wells correlated negatively with RDA1. The second axis (RDA2) separated mainly the background well from the rest of the wells located in the contaminated aquifer, but also separated the proximal well from the distal well.

The groundwater physicochemical parameters showed two major gradients: those that showed higher levels towards the proximal well




Fig. 6. RDA of microbial composition using the full OTU dataset among the sampling wells: R0 (background well), R1 (proximal well), R2 (intermediate well) and R4 (distal well). For clarity of readability, only OTUs with prominent vector length are shown.

and those that showed higher levels towards the distal well. The former included pH, ammonium, calcium, alkalinity, sulphate, conductivity, sodium, chloride and potassium, which have correlated negatively with RDA2. The second gradient was driven by dissolved oxygen, magnesium, iron and manganese, which have likewise correlated negatively with RDA2, except dissolved oxygen that correlated positively with RDA2. Nitrate and magnesium have shown moderate gradient between the proximal and distal wells, while TOC showed higher levels towards the intermediate well.

The prominent OTUs in the intermediate and background wells were dominated by uncultured lineages. On the other hand, the abundant OTUs in the distal and proximal wells composed mainly culturable taxa, although those in the proximal well seem to constitute those requiring special growth conditions. Some of the OTUs that showed prominent vectors in the RDA analysis were also among the abundant (>2%) OTUs. They include *Alkanindiges*, *Duganella*, *Undibacterium*, GKS98 (uncultured *Alcaligenaceae*), *Sulfuritalea*, *Sulfurifustis*, *Thermomonas*, *Rhodanobacter*, *Brevundimonas*, *Methylobacter*, DTB120 and UBA12409 (uncultured *Babeliales*).

3.2.5. Predicted phenotypic functions

Taxonomic-to-phenotypic mapping using METAGENassist (Fig. 7) predicted 15 potential energy source phenotypes, predominantly phototroph, heterotroph, autotroph and organotroph, methanotroph, methylotroph and oligotroph, in varying abundances. Five groups were present at a relative abundance >0.2. These were phototroph in the background and distal wells, heterotroph in the background and proximal wells, organotroph in the proximal well, and autotroph and methanotroph in the intermediate well. Heterotroph and phototroph accounted for >50% of the total abundance in the background and distal wells, respectively. Overall, the background well consisted of two major energy source-types, the proximal and intermediate wells seven and six, respectively, and the distal well four.

Twenty-five different potential metabolic profiles were predicted (Fig. 8). Degradors of both inorganic and organic compounds were predicted. Organic compound transformers were, however, in lower abundances than inorganic compound metabolisers. Microbes involved in sulphur and nitrogen transformations formed the dominant groups. Generally, the abundance of metabolic profiles varied from well to well.

4. Discussion

4.1. Cell density and microbial diversity

Overall, cell density estimate from microscopic count was significantly higher than plate count, which is consistent with the literature (Gregorich and Carter, 2007; Muyzer and Smalla, 1998) and expected from theoretical considerations. However, within the wells located in the contaminated aquifer, this difference was seldom greater than a factor of two in most cases, and only in three of the twelve multilevel samples was the difference significant. This suggests that in most samples, a large proportion of the microbial population is culturable. The compounds in the landfill leachate likely favoured the growth of culturable heterotrophic microorganisms. The physiological profiles of microbial communities in aquifers receiving a landfill leachate are expected to be due to culturable bacteria (Röling et al., 2000). However, the metabarcoding data conflict with these findings, the highest proportion of uncultured OTUs being found in the intermediate well, rather than in the distal well as indicated by the comparison of microscopic and plate counts.

The background aquifer had low solute levels, indicating that the aquifer is nutrient-poor. Based on the culture method, both cell density and the microbial diversity were low, although a comparison of microscopic and plate counts suggests that a fairly large proportion of the population is non-culturable. This is consistent with the metabarcoding data, in which the microbiome of the background aquifer was dominated by uncultured taxa and only a few culturable taxa such as *Duganella*, *Rhodospirillum rubrum* and *Alkanindiges* were abundant. Similarly, the Shannon diversity index (3.57 ± 1.32) was lower than the overall diversity index recorded in the contaminated aquifer (3.96 ± 0.71). Further, the RDA ordination (Fig. 6) indicates that the background well is compositionally very different from the contaminated aquifer. Similar differences in microbial communities between contaminated and uncontaminated groundwater have been reported (Brad et al., 2013; Brad et al., 2008; Mouser et al., 2005).

The contaminated aquifer, on the other hand, is relatively solute-rich and supports a denser and more diverse microbial population; a scenario likely to occur where a landfill leaches easily degradable organic matter (Röling et al., 2000). Although the concentrations of solutes




Fig. 7. Predicted energy source requirements of the OTUs in R0, R1, R2 and R4. R0 is the background well located in a nearby uncontaminated aquifer, while R1, R2 and R4 are the wells located in the contaminated aquifer placed at the proximal, intermediate and distal positions from the landfill. The vertical dashed line depicts 20% abundance.




Fig. 8. Predicted metabolic potentials of the OTUs in R0, R1, R2 and R4. R0 is the background well located in a nearby uncontaminated aquifer, while R1, R2 and R4 are the wells located in the contaminated aquifer placed at the proximal, intermediate and distal positions from the landfill.

have decreased greatly compared to previous data (Abiriga et al., 2020), they are evidently still sufficient to maintain a distinctive bacterial community; a previous study (Herzyk et al., 2017) indicates that physicochemical parameters return to normal more quickly than the microbial community. In line with this, data from culturing, fluorescence microscopy and metabarcoding showed higher species and cell density, diversity and species richness in the contaminated aquifer. The highest species richness and Shannon entropy were recorded in the intermediate well (Fig. 5A-B and Table S4), an observation that agrees with the culture and microscopic methods, where the highest cell and plate counts were recorded in the intermediate well (Fig. 2). On the other hand, moderate and lowest species richness and diversity were recorded in the proximal and distal well, respectively (Fig. 5A-B). This suggests the existence of an ecological gradient along the groundwater flow path, in which the proximal and intermediate wells are expected to have a high resemblance, as they are spatially close to each other. This is consistent with both the culture and metabarcoding data (Fig. 3 and Fig. 5C), where the highest shared species and genera was recorded between the proximal and intermediate wells. Further, these wells clustered next to one another (Fig. 6), suggesting that they are relatively similar and are thus made up of fairly similar microbial communities. The distal well is therefore considered to be ecologically more different from the upstream wells, which is consistent with the observations that the highest unique species (28) (Fig. 3B) and genera (90) (Fig. 5C) were found in the distal well. This is further supported by the beta diversity (Fig. 6) which indicates that the microbial composition of the distal well is different from the upstream wells.

The concentration of some of the groundwater physicochemical variables was found to decrease with distance from the landfill (Table 1),

and this was accompanied by an increase in microbial diversity (Fig. 5 and Table S4) at least from the proximal to intermediate wells. This may be analogous to increase in diversity and community stability as leachate becomes less contaminated over time (Köchling et al., 2015). Thus, close to the landfill, only species resistant to the toxic effects of the leachate are probably able to survive and grow, while as toxic pollutants become attenuated through biotic and abiotic processes, it allows the growth of more sensitive species. A previous study from Norman Landfill in the United States has shown that microbial gene diversity varied with distance from the landfill (Lu et al., 2012). In either study (both the previous and the present), the observed changes in diversity as a function of distance indicates the significance of attenuation mechanisms *in situ* that shape the microbial composition and function. Such spatial variation may enhance biodegradation as pollutants migrate from one region with a given microbial composition to another of a different microbial composition (Brad et al., 2013; Brad et al., 2008; Mouser et al., 2005; Röling et al., 2000).

4.2. Microbial compositions and environmental significance

Four microbial phyla were identified by the culture method (Fig. S3) compare to fifty-seven identified by the metabarcoding of which only eight occurred in greater (>3.5%) abundances. Of these eight phyla, *Proteobacteria*, *Bacteroidetes*, *Actinobacteria* and *Firmicutes* were represented among the cultured isolates, while *Patensibacteria*, *Chloroflexi*, *Acidobacteria* and *Verrucomicrobia* were only present in the metabarcoding data. *Proteobacteria* was the most abundant phylum and accounted for 31% and 55.2% of the total microbiome in the metabarcoding and culture data, respectively. *Bacteroidetes* (16.1%)

and *Actinobacteria* (25.3%) were present in comparable proportions, while *Firmicutes* (3.4%) was the least abundant of the eight phyla (based on culture data).

Putative hydrocarbon degraders among the isolates are presented in Table S5. Those from the metabarcoding data include the antipyrine/chloridazon-degrader *Phenylobacterium* (Lingens et al., 1985; Oh and Roh, 2012), alkane/alkyl-degraders *Parvibaculum* and *Alkanindiges* (Bogan et al., 2003; Lai et al., 2011; Schleheck et al., 2004), ibuprofen-degrader *Patulibacter medicamentivorans* (Almeida et al., 2013) and oil-degrader *Aquabacterium* (*A. olei*) (Jeong and Kim, 2015). Others include the single-carbon metabolisers *Methylobacter*, *Methylotenera* and *Methylocystis*. Energy source prediction showed the presence of methylotrophs and methanotrophs in high proportions in the contaminated wells (R1-R4) (Fig. 7), which may suggest the presence of methane in the aquifer. The landfill has now matured (manuscript under consideration) and is supposedly in transition from methanogenic to aerobic stage, i.e., from phase III to phase IV of landfill stabilisation (Kjeldsen et al., 2002), but there could still be active methanogenesis. A wide range of hydrocarbon metabolisms were also predicted through the phenotypic mapping (Fig. 8). Although the TOC in the groundwater was low with a weak gradient along the groundwater flow path, it could still provide carbon source to the aquifer microbiome. Considering the age of the landfill, the TOC should be predominantly of recalcitrant fractions (Kulikowska and Klimiuk, 2008) and this makes it difficult to degrade (Appelo and Postma, 2005), leading to a non-significant difference across the wells.

From the metabarcoding data, sulphur cyclers were detected and included the sulphur/sulphide oxidisers *Sulfuritalea* and *Sulfurifustis* (Kojima et al., 2015; Kojima and Fukui, 2011) (Fig. 4), and *Sulfuricella* (Kojima and Fukui, 2010) and *Rhodobacter* (Imhoff et al., 1984) (Fig. 6). However, sulphate showed a weak gradient with a non-significant difference across the wells (Table S1), suggesting that the aquifer contains a significant sulphate sink, such as re-reduction in local anaerobic pockets.

Both nitrate reducers and ammonia oxidisers were identified. Nitrate reducers included 29 cultured isolates (Table S6), and genera *Cavicella*, *Sterolibacterium*, *Aquabacterium* and *Novosphingobium* from the metabarcoding data. *Sulfuricella*, which at present is monotypic (*S. denitrificans*), correlated positively with nitrate and could therefore be involved in denitrification. Among the nitrogen oxidisers were the ammonia oxidisers *Nitrosospora* (Watson, 1971) and nitrite oxidisers *Nitrospinaeae* (Lücker and Daims, 2014). Taxonomic-to-phenotypic mapping (Fig. 8) indicates the presence of ammonia oxidisers and nitrate/nitrite reducers. Both ammonium and nitrate were detectable in the groundwater, although nitrate was present in higher levels than ammonium. The presence of both ions may suggest complete cycling of nitrogen, although the reduction reaction might be limited due to low organic matter. The lack of a significant difference in nitrate across the wells suggests a balance between reductive and oxidative nitrate metabolism.

Iron metabolisers include *Gallionella*, a genus known for centuries to clog well screens (Chapelle, 2001) by oxidising ferrous iron at an ecotone between reducing and oxidising environments. Genus *Rhodospirillum*, which the culture-based analysis showed to be represented by *R. ferrireducens*, together with genus *Ferribacterium*, are both iron-reducing bacteria; a role that counteracts that played by *Gallionella*. In addition, *R. ferrireducens* is known to carry out metabolism of manganese and oxygen (Finneran et al., 2003). Although the level of iron was low across the wells, the distal well where *Rhodospirillum* and *Ferribacterium* were abundant was slightly enriched in iron, manganese and oxygen (Table 1 and Fig. 6). The greater abundance of iron-reducers in the background well (Fig. 8) can be attributed to *Rhodospirillum*, although iron in this well was below the limit of detection. Like sulphate reduction, iron reduction must be limited due to the low organic matter. We propose that the high proportion of phototrophs (Fig. 7) is due to the high abundance of *Rhodospirillum* found in the respective wells since two species of this genus are phototrophs (Finneran et al., 2003).

Although phototrophy does not mean being obligately photosynthetic, the result here is likely due to misclassification as the programme (METAGENassist) uses OTU information which in metabarcoding are seldom resolved to species-level and unambiguous functional assignment is, therefore, currently a methodological limitation.

Genus *Polynucleobacter* was one of the abundant taxa (Fig. 4). Members of the genus are obligate endosymbionts of ciliates (Heckmann and Schmidt, 1987), and thus the presence of a substantial protistan population is indicated, which is further supported by the presence of *Legionella*, a parasite of amoebae. In addition, *Polynucleobacter*, and other genera such as *Aquabacterium*, *Rhodospirillum*, *Duganella* and *Limnochlamydomonas*, were disproportionately more abundant in spring (Fig. S6). The seasonal variation in the microbial community composition will be addressed in a future manuscript (under consideration). *Duganella* showed higher abundance in the background and distal wells (Fig. 4). The genus showed stronger correlation with dissolved oxygen (Fig. 6) and given that dissolved oxygen was higher in these wells, it might be a relevant selection factor for the genus.

5. Conclusion

This study reports the microbial diversity of a landfill-contaminated confined aquifer. Both culture-dependent and culture-independent microbial techniques were applied. Comparisons of microscopic cell counts and plate counts as well as metabarcoding data suggest that the microbes in the most contaminated part of the aquifer were mostly culturable heterotrophs. In the contaminated aquifer, microbial diversity was moderate in the proximal well, highest in the intermediate well and lowest in the distal well. The lower diversity in the distal well indicates dominance of the microbial community by a few taxa. Each well had a distinctive microbial flora, but the proximal and intermediate wells seemed to be ecologically related as they had more similar microbial community composition. Comparison between the uncontaminated and contaminated aquifers showed higher microbial diversity in the contaminated aquifer. Compositionally, there was a clear difference between the flora of the contaminated wells and the uncontaminated well. The data suggests that the microbiome of the contaminated aquifer has been impacted by the landfill leachate. Functional analysis indicates the presence of microbes capable of hydrocarbon, sulphur, nitrogen, iron and manganese metabolism.

CRedit authorship contribution statement

Daniel Abiriga: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – original draft, Writing – review & editing. **Andrew Jenkins:** Conceptualization, Methodology, Supervision, Writing – review & editing. **Kristian Alfnes:** Formal analysis, Software, Validation, Writing – review & editing. **Live S. Vestgarden:** Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Supervision, Writing – review & editing. **Harald Klempe:** Conceptualization, Methodology, Project administration, Resources, Supervision, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

We thank Frode Bergan and Tom Aage Aarnes for participation in fieldwork, Karin Brekke Li for technical assistance in water chemistry analysis and Benedikte N. Pedersen for technical advice during sequencing of bacterial isolates. Fluorescence microscopy was conducted at the

department of process, energy and environment, University of South-Eastern Norway and we thank the late Professor Rune Bakke for providing access to the instrument and Ikumi Umetani for help with its use. The sequencing service was provided by the Norwegian Sequencing Centre (<https://www.sequencing.uio.no/>), a national technology platform hosted by the University of Oslo and supported by the Functional Genomics and Infrastructure programs of the Research Council of Norway and the Southeastern Regional Health Authorities. We thank the two anonymous reviewers for their comments which improved the quality of the manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2021.147349>.

References

- Abiriga, D., Vestgarden, L.S., Klempe, H., 2020. Groundwater contamination from a municipal landfill: effect of age, landfill closure, and season on groundwater chemistry. *Sci. Total Environ.* 737, 140307.
- Abiriga, D., Vestgarden, L.S., Klempe, H., 2021. Long-term redox conditions in a landfill-leachate-contaminated groundwater. *Sci. Total Environ.* 755, 143725.
- Albrechtsen, H.-J., Heron, G., Christensen, T.H., 1995. Limiting factors for microbial Fe (III)-reduction in a landfill leachate polluted aquifer (Vejen, Denmark). *FEMS Microbiol. Ecol.* 16 (3), 233–247.
- Alfreider, A., Vogt, C., Babel, W., 2002. Microbial diversity in an in situ reactor system treating onchlorobenzene contaminated groundwater as revealed by 16S ribosomal DNA analysis. *Syst. Appl. Microbiol.* 25 (2), 232–240.
- Almeida, B., Vaz-Moreira, I., Schumann, P., Nunes, O.C., Carvalho, G., Crespo, M.T.B., 2013. *Patulibacter medicamentivorans* sp. nov., isolated from activated sludge of a wastewater treatment plant. *Int. J. Syst. Evol. Microbiol.* 63 (7), 2588–2593.
- Anantharaman, K., Brown, C.T., Hug, L.A., Sharon, I., Castelle, C.J., Probst, A.J., Thomas, B.C., Singh, A., Wilkins, M.J., Karaoz, U., 2016. Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system. *Nat. Commun.* 7.
- Anderson, M.J., 2006. Distance-based tests for homogeneity of multivariate dispersions. *Biometrics* 62 (1), 245–253.
- Appelo, C.A.J., Postma, D., 2005. *Geochemistry, Groundwater and Pollution* (Balkema, Leiden).
- Arndt, D., Xia, J., Liu, Y., Zhou, Y., Guo, A.C., Cruz, J.A., Sinelnikov, I., Budwill, K., Nesbø, C.L., Wishart, D.S., 2012. METAGENassist: a comprehensive web server for comparative metagenomics. *Nucleic Acids Res.* 40(W1), W88–W95.
- Bogan, B.W., Sullivan, W.R., Kayser, K.J., Derr, K., Aldrich, H.C., Paterek, J.R., 2003. *Alkanindiges illinoisensis* gen. nov., sp. nov., an obligately hydrocarbonoclastic, aerobic squalene-degrading bacterium isolated from oilfield soils. *J. Med. Microbiol.* 53 (5), 1389–1395.
- Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* 37 (8), 852–857.
- Brad, T., Van Breukelen, B.M., Braster, M., Van Straalen, N.M., Röling, W.F., 2008. Spatial heterogeneity in sediment-associated bacterial and eukaryotic communities in a landfill leachate-contaminated aquifer. *FEMS Microbiol. Ecol.* 65 (3), 534–543.
- Brad, T., Oberfell, C., van Breukelen, B., van Straalen, N., Röling, W., 2013. Spatiotemporal variations in microbial communities in a landfill leachate plume. *Groundwater Monitoring & Remediation* 33 (4), 69–78.
- Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13 (7), 581.
- Chapelle, F.H., 2001. *Ground-Water Microbiology and Geochemistry*. Wiley, New York.
- Chen, Q.-L., Li, H., Zhou, X.-Y., Zhao, Y., Su, J.-Q., Zhang, X., Huang, F.-Y., 2017. An underappreciated hotspot of antibiotic resistance: the groundwater near the municipal solid waste landfill. *Sci. Total Environ.* 609, 966–973.
- Christensen, T.H., Bjerg, P.L., Kjeldsen, P., 2000. Natural attenuation: a feasible approach to remediation of ground water pollution at landfills? *Groundwater Monitoring & Remediation* 20 (1), 69–77.
- Cozzarelli, I.M., Sufflita, J.M., Ulrich, G.A., Harris, S.H., Scholl, M.A., Schlottmann, J.L., Christenson, S., 2000. Geochemical and microbiological methods for evaluating anaerobic processes in an aquifer contaminated by landfill leachate. *Environ. Sci. Technol.* 34 (18), 4025–4033.
- Csuros, M., Csuros, C., Ver, K., 1999. *Microbiological Examination of Water and Wastewater*. Lewis Publishers, Boca Raton, Fla.
- Dlugonski, J., 2016. *Microbial Biodegradation: From Omics to Function and Application*. Caister Academic Press, Norfolk.
- Dojka, M.A., Hugenholtz, P., Haack, S.K., Pace, N.R., 1998. Microbial diversity in a hydrocarbon- and chlorinated-solvent-contaminated aquifer undergoing intrinsic bioremediation. *Appl. Environ. Microbiol.* 64 (10), 3869–3877.
- Eggen, T., Moeder, M., Arukwe, A., 2010. Municipal landfill leachates: a significant source for new and emerging pollutants. *Sci. Total Environ.* 408 (21), 5147–5157.
- Fadrosch, D.W., Ma, B., Gajer, P., Sengamaly, N., Ott, S., Brotman, R.M., Ravel, J., 2014. An improved dual-indexing approach for multiplexed 16S rRNA gene sequencing on the Illumina MiSeq platform. *Microbiome* 2 (1), 6.
- Finneran, K.T., Johnsen, C.V., Lovley, D.R., 2003. *Rhodoferrax ferrireducens* sp. nov., a psychrotolerant, facultatively anaerobic bacterium that oxidizes acetate with the reduction of Fe (III). *Int. J. Syst. Evol. Microbiol.* 53 (3), 669–673.
- Gregorich, E.G., Carter, M.R., 2007. *Soil Sampling and Methods of Analysis*. CRC Press.
- Harvey, R.W., Smith, R.L., George, L., 1984. Effect of organic contamination upon microbial distributions and heterotrophic uptake in a Cape Cod, Mass., aquifer. *Appl. Environ. Microbiol.* 48 (6), 1197–1202.
- He, Z., Zhang, P., Wu, L., Rocha, A.M., Tu, Q., Shi, Z., Wu, B., Qin, Y., Wang, J., Yan, Q., 2018. Microbial functional gene diversity predicts groundwater contamination and ecosystem functioning. *MBio* 9 (1), e02417–e02425.
- Heckmann, K., Schmidt, H.J., 1987. *Polynucleobacter necessarius* gen. nov., sp. nov., an obligately endosymbiotic bacterium living in the cytoplasm of *Euplotes aediculatus*. *Int. J. Syst. Evol. Microbiol.* 37 (4), 456–457.
- Herzyk, A., Fillingner, L., Larentis, M., Qiu, S., Malozzewski, P., Hünninger, M., Schmidt, S.I., Stumpp, C., Marozava, S., Knappett, P.S., 2017. Response and recovery of a pristine groundwater ecosystem impacted by toluene contamination—a meso-scale indoor aquifer experiment. *J. Contam. Hydrol.* 207, 17–30.
- Hug, L.A., Thomas, B.C., Brown, C.T., Frischkorn, K.R., Williams, K.H., Tringe, S.G., Banfield, J.F., 2015. Aquifer environment selects for microbial species cohorts in sediment and groundwater. *The ISME Journal* 9 (8), 1846–1856.
- Hyldegaard, B.H., Jakobsen, R., Weeth, E.B., Overheu, N.D., Gent, D.B., Ottosen, L.M., 2019. Challenges in electrochemical remediation of chlorinated solvents in natural groundwater aquifer settings. *J. Hazard. Mater.* 368, 680–688.
- Imhoff, J., Trüper, H., Pfennig, N., 1984. Rearrangement of the species and genera of the phototrophic "purple nonsulfur bacteria". *Int. J. Syst. Evol. Microbiol.* 34 (3), 340–343.
- Jeong, S.-W., Kim, J., 2015. *Aquabacterium olei* sp. nov., an oil-degrading bacterium isolated from oil-contaminated soil. *Int. J. Syst. Evol. Microbiol.* 65 (10), 3597–3602.
- Kepner, R., Pratt, J.R., 1994. Use of fluorochromes for direct enumeration of total bacteria in environmental samples: past and present. *Microbiol. Mol. Biol. Rev.* 58 (4), 603–615.
- Kjeldsen, P., Barlaz, M.A., Rooker, A.P., Baun, A., Ledin, A., Christensen, T.H., 2002. Present and long-term composition of MSW landfill leachate: a review. *Crit. Rev. Environ. Sci. Technol.* 32 (4), 297–336.
- Kleikemper, J., Pombo, S.A., Schroth, M.H., Sigler, W.V., Pesaro, M., Zeyer, J., 2005. Activity and diversity of methanogens in a petroleum hydrocarbon-contaminated aquifer. *Appl. Environ. Microbiol.* 71 (1), 149–158.
- Klempe, H., 2004. Identification of Quaternary subsurface glacial deposits using 3D databases and GIS. *Norsk Geografisk Tidsskrift-Norwegian Journal of Geography* 58 (2), 90–95.
- Klempe, H., 2015. The hydrogeological and cultural background for two sacred springs, Bø, Telemark County, Norway. *Quat. Int.* 368, 31–42.
- Köchling, T., Sanz, J.L., Gavazza, S., Florencio, L., 2015. Analysis of microbial community structure and composition in leachates from a young landfill by 454 pyrosequencing. *Appl. Microbiol. Biotechnol.* 99 (13), 5657–5668.
- Kojima, H., Fukui, M., 2010. *Sulfuricella denitrificans* gen. nov., sp. nov., a sulfur-oxidizing autotroph isolated from a freshwater lake. *Int. J. Syst. Evol. Microbiol.* 60 (12), 2862–2866.
- Kojima, H., Fukui, M., 2011. *Sulfuritalea hydrogenivorans* gen. nov., sp. nov., a facultative autotroph isolated from a freshwater lake. *Int. J. Syst. Evol. Microbiol.* 61 (7), 1651–1655.
- Kojima, H., Shinohara, A., Fukui, M., 2015. *Sulfurifustis variabilis* gen. nov., sp. nov., a sulfur oxidizer isolated from a lake, and proposal of *Acidiferrobacteraceae* fam. nov. and *Acidiferrobacterales* ord. nov. *Int. J. Syst. Evol. Microbiol.* 65 (10), 3709–3713.
- Kulikowska, D., Klimiuk, E., 2008. The effect of landfill age on municipal leachate composition. *Bioresour. Technol.* 99 (13), 5981–5985.
- Lai, Q., Wang, L., Liu, Y., Yuan, J., Sun, F., Shao, Z., 2011. *Parvibaculum indicum* sp. nov., isolated from deep-sea water. *Int. J. Syst. Evol. Microbiol.* 61 (2), 271–274.
- Legendre, P., Gallagher, E.D., 2001. Ecologically meaningful transformations for ordination of species data. *Oecologia*. vol. 129(2), pp. 271–280.
- Lin, B., Braster, M., Röling, W.F., van Breukelen, B.M., 2007. Iron-reducing microorganisms in a landfill leachate-polluted aquifer: complementing culture-independent information with enrichments and isolations. *Geomicrobiol. J.* 24 (3–4), 283–294.
- Lingens, F., Blecher, R., Blecher, H., Blobel, F., Eberspächer, J., Fröhner, C., Görisch, H., Görisch, H., Layh, G., 1985. *Phenylobacterium immobile* gen. nov., sp. nov., a gram-negative bacterium that degrades the herbicide chloridazon. *Int. J. Syst. Evol. Microbiol.* 35 (1), 26–39.
- Logeswaran, P., Megharaj, M., Chadalavada, S., Bowman, M., Naidu, R., 2018. Petroleum hydrocarbons (PH) in groundwater aquifers: an overview of environmental fate, toxicity, microbial degradation and risk-based remediation approaches. *Environ. Technol. Innov.* 10, 175–193.
- Lu, Z., He, Z., Parisi, V.A., Kang, S., Deng, Y., Van Nostrand, J.D., Masoner, J.R., Cozzarelli, I.M., Sufflita, J.M., Zhou, J., 2012. GeoChip-based analysis of microbial functional gene diversity in a landfill leachate-contaminated aquifer. *Environ. Sci. Technol.* 46 (11), 5824–5833.
- Lücker, S., Daims, H., 2014. In: Rosenberger, E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F. (Eds.), *The Prokaryotes: Deltaproteobacteria and Epsilonproteobacteria*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 231–237.
- Ludvigsen, L., Albrechtsen, H.-J., Ringelberg, D., Ekelund, F., Christensen, T., 1999. Distribution and composition of microbial populations in a landfill leachate contaminated aquifer (Grindsted, Denmark). *Microb. Ecol.* 37 (3), 197–207.
- Mays, D., Scheibe, T., 2018. *Groundwater Contamination, Subsurface Processes, and Remediation Methods: Overview of the Special Issue of Water on Groundwater Contamination and Remediation*. Multidisciplinary Digital Publishing Institute.
- Meckenstock, R.U., Elsner, M., Griebler, C., Lueders, T., Stumpp, C., Aamand, J., Agathos, S.N., Albrechtsen, H.-J., Bastiaens, L., Bjerg, P.L., 2015. Biodegradation: Updating the

- Concepts of Control for Microbial Cleanup in Contaminated Aquifers. ACS Publications.
- Mouser, P.J., Rizzo, D.M., Röling, W.F., van Breukelen, B.M., 2005. A multivariate statistical approach to spatial representation of groundwater contamination using hydrochemistry and microbial community profiles. *Environ. Sci. Technol.* 39 (19), 7551–7559.
- Muyzer, G., Smalla, K., 1998. Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Antonie Van Leeuwenhoek* 73 (1), 127–141.
- Nunes, O.C., Lopes, A.R., Manaia, C.M., 2013. Microbial degradation of the herbicide molinate by defined cultures and in the environment. *Appl. Microbiol. Biotechnol.* 97 (24), 10275–10291.
- O'Connor, D., Hou, D., Ok, Y.S., Song, Y., Sarmah, A.K., Li, X., Tack, F.M., 2018. Sustainable in situ remediation of recalcitrant organic pollutants in groundwater with controlled release materials: a review. *J. Control. Release* 283, 200–213.
- Oh, Y.-S., Roh, D.-H., 2012. *Phenylobacterium muchangponense* sp. nov., isolated from beach soil, and emended description of the genus *Phenylobacterium*. *Int. J. Syst. Evol. Microbiol.* 62 (4), 977–983.
- Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. *vegan: Community Ecology Package* <https://CRAN.R-project.org/package=vegan>.
- Pickup, R., Rhodes, G., Alamillo, M., Mallinson, H., Thornton, S., Lerner, D., 2001. Microbiological analysis of multi-level borehole samples from a contaminated groundwater system. *J. Contam. Hydrol.* 53 (3–4), 269–284.
- Pilloni, G., Bayer, A., Ruth-Anneser, B., Fillinger, L., Engel, M., Griebler, C., Lueders, T., 2019. Dynamics of hydrology and anaerobic hydrocarbon degrader communities in a tar-oil contaminated aquifer. *Microorganisms* 7 (2), 46.
- Pous, N., Balaguer, M.D., Colprim, J., Puig, S., 2018. Opportunities for groundwater microbial electro-remediation. *Microb. Biotechnol.* 11 (1), 119–135.
- R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing <https://www.R-project.org/> (Vienna, Austria).
- Reinhard, M., Barker, J.F., Goodman, N.L., 1984. Occurrence and distribution of organic chemicals in two landfill leachate plumes. *Environ. Sci. Technol.* 18 (12), 953–961.
- Röling, W., Van Breukelen, B., Braster, M., Goeltom, M., Groen, J., Van Verseveld, H., 2000. Analysis of microbial communities in a landfill leachate polluted aquifer using a new method for anaerobic physiological profiling and 16S rDNA based fingerprinting. *Microb. Ecol.* 40 (3), 177–188.
- Röling, W.F., van Breukelen, B.M., Braster, M., Lin, B., van Verseveld, H.W., 2001. Relationships between microbial community structure and hydrochemistry in a landfill leachate-polluted aquifer. *Appl. Environ. Microbiol.* 67 (10), 4619–4629.
- Rooney-Varga, J.N., Anderson, R.T., Fraga, J.L., Ringelberg, D., Lovley, D.R., 1999. Microbial communities associated with anaerobic benzene degradation in a petroleum-contaminated aquifer. *Appl. Environ. Microbiol.* 65 (7), 3056–3063.
- Schleheck, D., Tindall, B.J., Rossello-Mora, R., Cook, A.M., 2004. *Parvibaculum lavamentivorans* gen. nov., sp. nov., a novel heterotroph that initiates catabolism of linear alkylbenzenesulfonate. *Int. J. Syst. Evol. Microbiol.* 54 (5), 1489–1497.
- Taş, N., Brandt, B.W., Braster, M., van Breukelen, B.M., Röling, W.F., 2018. Subsurface landfill leachate contamination affects microbial metabolic potential and gene expression in the Banisveld aquifer. *FEMS Microbiol. Ecol.* 94 (10), fty156.
- Tischer, K., Zeder, M., Klug, R., Pernthaler, J., Schattenhofer, M., Harms, H., Wendeberg, A., 2012. Fluorescence in situ hybridization (CARD-FISH) of microorganisms in hydrocarbon contaminated aquifer sediment samples. *Syst. Appl. Microbiol.* 35 (8), 526–532.
- Watanabe, K., Kodama, Y., Hamamura, N., Kaku, N., 2002. Diversity, abundance, and activity of archaeal populations in oil-contaminated groundwater accumulated at the bottom of an underground crude oil storage cavity. *Appl. Environ. Microbiol.* 68 (8), 3899–3907.
- Watson, S.W., 1971. Taxonomic considerations of the family Nitrobacteraceae Buchanan requests for Opinions 1. *Int. J. Syst. Evol. Microbiol.* 21 (3), 254–270.