

Population and distribution of European beavers (*Castor fiber*)

Duncan J. Halley¹ & Frank Rosell²

¹Norwegian Institute for Nature Research, Tungasletta 2, N-7485 Trondheim, Norway,
e-mail: duncan.halley@nina.no

²Telemark University College, N-3800 Bø i Telemark, Norway

Abstract: After being reduced to about 1,200 animals in eight isolated populations by the beginning of the 20th century, European beavers (*Castor fiber*) have powerfully recovered in both range and population, through relaxation of persecution, natural spread, and widespread reintroductions. Populations are now (2003) established in all countries within their former natural range in Europe except for Britain, Portugal, Italy, and the south Balkans (Greece, Albania, Bulgaria, Macedonia; status in Bosnia-Herzegovina is uncertain). In Asia, there are significant populations in central Siberia, Kamchatka, and on the Amur; and small relict populations elsewhere in Siberia, and in Xinjiang (China)/western Mongolia. The current minimum population estimate is 639,000. Both populations and range are in rapid expansion. We present maps summarizing current knowledge of the world distribution of European beaver and the Eurasian distribution of the introduced American beaver (*Castor canadensis*), and tables of the most recent known population estimates for each country.

Keywords: beaver, *Castor fiber*, population, distribution, range, reintroduction.

Introduction

European beavers (*Castor fiber*) have, since the late 19th century, staged a remarkable recovery both in population and distribution. From being a critically endangered species reduced to about 1,200 animals in scattered enclaves, the species is now conservationally secure and, aided by widespread reintroductions, rapidly recolonising much of its range, including areas where it has not occurred for centuries or even millennia. The pattern of reexpansion is not only interesting in itself, but also offers valuable insights in the fields of population biology and conservation ecology. Until recently, this expansion has not been well documented, but since the latter part of the 20th century increasing amounts of information have become available, allowing the preparation of reviews summarizing the then current status of the species (Macdonald et al. 1995, Halley & Rosell 2002). However, both popula-

tion and distribution continue to expand rapidly, both through natural spread and new reintroductions, so that these reviews rapidly become overtaken by events. Regular summaries of the most recent population and distribution data are therefore of use, both to current researchers and managers seeking an overview of the many, widely scattered, papers on the subject, and for future biologists interested in following in detail a remarkable case study in conservation biology.

Distribution

Population distribution is best known in western and central Europe (figure 1), less so for European Russia and Asia (figure 2). The continuous population ranges from eastern Poland through the Baltic States and European Russia to central Siberia. There is a large disjunct population in Norway and Sweden, and smaller scattered disjunct populations through the rest of mainland Europe. Disjunct reintroduced populations are also found on the periphery of the main Russian range, on the Amur watershed in eastern Siberia,




Figure 1. Distribution of beavers in Europe, excluding Russia. Locations of relict populations are marked in black. Traditional subspecies designations: 1 = *Castor fiber fiber*; 2 = *Castor fiber albicus*; 3 = *Castor fiber galliae*; 4 = *Castor fiber belarusicus*. 1-3 are all *Castor fiber fiber*, 4 is *Castor fiber vistulanus* in DuCroz's reclassification (J.-F. DuCroz, personal communication; see Discussion). Dark shading represents the present range of *Castor fiber*; light shading represents the range of *Castor canadensis* in Finland. Squares are reintroduction sites where range has not yet spread significantly; crosses represent planned reintroductions (sources: Andersen 2002, P. Asbirk, personal communication, Balodis 1994, O. Boszér, personal communication, Bevanger 1995, O. Capt, personal communication, A. Czech, personal communication, Danilov 1995, Dúha & Majzlan 1997, Ermala et al. 1999, Fustec et al 2001, M. Grubestic, personal communication, Hartman 1999, G. Hartman, personal communication, Heideke & Ibe 1997, G. Ionescu, personal communication, Laanetu 1995, H. Lea, personal communication, V. Kostkan, personal communication, Mickus 1995, Niewold & Lammertsma 2000, F. Niewold, personal communication, Nolet & Rosell 1998, Office Nationale de la Chasse 1997, Pachinger & Hulik 1999, J. Ramon, personal communication, J. Reinhold, personal communication, Richard 1986, Rosell & Pedersen 1999, A.P. Saveljev, personal communication, G. Schwab, personal communication, J. Sieber, personal communication, H. Sluiter, personal communication, Troidl & Ionescu 1997, Ulevicius et al. 1999, A. Ulevicius, personal communication, D. Valachovic 1997, A. Vorel, personal communication, Winter 1997).

and on the Kamchatka peninsula in the Russian Far East. Although natural spread has contributed significantly to range and populations, most of the expansion is due to reintroductions, of which at least 157 have been recorded outside the former Soviet Union (FSU) (beaver were also extensively translocated within the FSU, but details are not available) (Halley & Rosell 2002). In 2003, beavers were reintroduced to the Ebro in Spain, with plans for further reintroductions to the Guadalquivir, Guadiana, and Tajo (Tagus) river systems (H. Lea, personal communication); reintroductions to Serbia at Obedska Bara and at Zasavica, on the Sava west of Belgrade, took place in spring 2004, and a later release to Ola Becej on the Danube north of Belgrade is planned (G. Schwab & D. Cirovic, personal communication). In Siberia there have been recent reintroductions (2001-2003) on the lower Ob and on the middle Lena (A.P. Saveljev, personal communication). Introduced populations of American beaver (*Castor canadensis*) are established in Finland and northwest Russia, on the Amur, and in Kamchatka (Saveljev & Safonov 1999, A.P. Saveljev, personal communication).

Available information on relict population sizes is given in table 1, and data on date of extinction, legal protection, reintroduction, and current population size in table 2.

Discussion

Populations of the eight refugia in which beavers survived the 19th century are currently each described as separate subspecies. However, recent mtDNA testing indicates clearly that only two subspecies are justifiable (*Castor fiber fiber* = *Castor fiber fiber*, *albicus* & *galliae*, and *Castor fiber vistulanus* = *Castor fiber belarusicus*, *osteuropaeus*, *pohlei*, *tuvanicus* & *birulei*) (J.-F. DuCroz, personal communication). This finding should be taken into account by managers seeking reintroduction stock in line with IUCN guidelines.

European beaver families consist of, on average, about five individuals, of which only the adult pair breeds. The minimum populations quoted should be interpreted in this light; the Rhône remnant population, for example, would seem to have been reduced to about six breeding pairs at minimum, well below the 25 pairs often quoted as a minimum viable population (MVP). (That evidence on the ground indicates that this rule-of-thumb is generally too pessimistic, is fortunate for conservation; see Caughley & Sinclair (1994) for review of this issue). The small size of remnant populations, and resultant inbreeding and loss of genetic diversity, does not seem to have led to breeding problems in this species, but




Figure 2. Distribution of beavers in Russia, Mongolia, Kazakhstan and Xinjiang (China). Locations of relict populations are marked in black. Traditional subspecies designations: 5 = *Castor fiber osteuropaeus*; 6 = *Castor fiber pohlei*; 7 = *Castor fiber tuvanicus*; 8 = *Castor fiber biruli*; 5-8 are all *Castor fiber vistulanus* in DuCroz's reclassification (J.-F. DuCroz, personal communication; see Discussion). Dark shading represents the present range of *Castor fiber*; light shading represents the range of *Castor canadensis* (figure adapted from Nolet & Rosell 1998 and A.P. Saveljev, personal communication).

Table 1. Location and estimated minimum population sizes of relict populations of European beaver (*Castor fiber*).

| Population | Minimum population size | References |
|--|-------------------------|-------------------------|
| Lower Rhône, France | 30 | Richard 1985 |
| Telemark, Norway | 60-120 | Collet 1897 |
| Elbe, Germany | 200 | Heideke & Hörig 1986 |
| Pripet marshes, Belarus/Ukraine/Russia | <300 | Zharkov & Solokov 1967 |
| Voronezh, Russia | 70 | Lavrov & Lavrov 1986 |
| Konda-Sosva, Russia | 300 | Lavrov & Lavrov 1986 |
| Upper Yenesei, Russia | 30-40 | Lavrov & Lavrov 1986 |
| Urungu, Mongolia/China | <100-150 | Lavrov & Hao-Tsuan 1961 |

suggests that populations descended solely from some of the smaller surviving populations may be more susceptible as a population to epidemic disease, due to lack of diversity in immune systems (Ellegren et al. 1993). In this connection, recent evidence that only two subspecies of European beaver can be justified (J.-F. DuCroz, personal communication; see above) suggests that managers may consider mixing individuals from the various refugia of each subspecies. This would broaden the genetic diversity within reintroduced populations somewhat without compromising subspecific integrity, as IUCN guidelines recommend.

The current world population estimate of 639,000 given here is conservative, calculated by adding together the lowest estimates for each country (rounded to the nearest thousand). All surviving European populations have grown in numbers beyond the point where further loss of genetic diversity through drift might be a conservation problem, though the status of some of the Siberian relict populations is unclear in this respect and would merit investigation.

The actual population is probably considerably higher than the figure given above. Assuming, for example, the higher end of the range estimates, and that populations estimated at 'greater than' were 10% above the figure given, yields a population estimate of just below 742,000. In some cases the population estimates given are several years old, so that, allowing in addition for natural increase, the true population may be closer to the higher of these figures.

However, the limitations in data available make all estimates beyond stating the minimum population speculative.

In most countries of the western and central European mainland, beavers exist at relatively low numbers for the present, and there is much unused suitable habitat. The range maps presented here are therefore probably conservative, as newly colonising beavers tend to select prime habitat requiring little alteration, where their presence may not immediately be noted by non-specialists. Very considerable expansion in both populations and range can be expected in the coming decades, especially in western Europe and the Danube watershed. A typical pattern of population development on a watershed following natural or artificial recolonisation has become evident in recent years, the evidence for which is reviewed in Halley & Rosell (2002); the most detailed case study of range and population expansion is found in Hartman (1995; personal communication.). At first, range expansion through the watershed is rapid, but population expansion is relatively slow. This seems to be because beavers select the best habitat available, rather than settling close to their natal territory. It is also difficult to find a mate in the vast, unoccupied stretches of a large watershed, and it appears that beaver will wander widely while searching. Extreme figures of movements of 500 km (Lavrov 1980 quoted in Saveljev et al. 2002), and 200 km involving crossing a watershed divide (Saveljev et al. 2002), have been reported, both of animals moving into uncolonised waters.

Table 2. The history and present status of European beavers (*Castor fiber*).

| Country | Extinction | Protection | Reintroduction and/or translocations | Present population size | References |
|----------------|-----------------------|------------|--------------------------------------|-------------------------|---|
| Austria | 1869 | – | 1970-1990 | >1300 | Kollar & Seiter 1990 Sieber 1999 J. Sieber, pers. comm. |
| Belarus | remnant | 1922 | – | 24,000 | Djoshkin & Safonov 1972 Safonov & Saveljev 1999 |
| Belgium | 1848 | – | 1998-2000 | 200-250 | Van den Bergh & Manet, pers. comm. |
| Bosnia-Herzeg. | ? | – | - ¹ | 0 ¹ | M. Grubestic, pers. comm. |
| Bulgaria | ? | – | planned | 0 | G. Schwab, pers. comm. |
| Croatia | 1857? | – | 1996-1998 | c.180 | M. Grubestic, pers. comm. |
| Czech Republic | 17th century | – | 1991-1992 1996 | c.500 | Kostkan & Lehky 1997 Kostkan 1999 |
| Denmark | c.500 BC ² | – | 1999 | 60-70 | Skov- og Naturstyrelsen 2000 Berthelsen & Madsen 2003 J. Berthelsen, pers. comm. |
| England | <12th century | – | – | 0 ³ | Macdonald et al. 1995 Campbell & Tattersall 2003 F. Tattersall, pers. comm. |
| Estonia | 1841 | – | 1957 | 11,000 | Laanetu 1995 U. Timm, pers. comm. A. Ulevicius, pers. comm. |
| Finland | 1868 | 1868 | 1935-1937 1995 | 2000 ⁴ | Lahti 1995 Ermala et al. 1999 A. Ermala, pers. comm. |
| France | remnant | 1909 | 1959-95 | 7000-10,000 | Richard 1985, 1986 Fustec et al. 2001 R. Dennis, pers. comm. P. Rouland, pers. comm. |
| Germany | remnant | 1910 | 1936-1940 1966-1989 1999-2000 | 8000-10,000 | Schwab et al. 1994 Macdonald et al. 1995 G. Schwab, pers. comm. |
| Hungary | 1865 | – | 1991-1993 1996-2003 | >400 | Kollar & Seiter 1990 O. Bozsér, pers. comm. G. Schwab, pers. comm. |

Table 2 (continued). The history and present status of European beavers (*Castor fiber*).

| Country | Extinction | Protection | Reintroduction and/or translocations | Present population size | References |
|----------------|---------------|------------|--------------------------------------|-------------------------|--|
| Italy | 1541 | – | proposed | 0 | Nolet 1996 |
| Kazakhstan | ? | – | – | 1000 | Djoshkin & Safonov 1972 Safonov & Saveljev 1999 Saveljev & Safonov 1999 |
| Latvia | 1830s | – | 1927-1952 1975-1984 | >100,000 | Balodis 1992, 1995, 1997, 1998 Ozolins & Baumanis 2000 |
| Lithuania | 1938 | – | 1947-1959 | 50,000- 70,000 | Palionene 1965 Mickus 1995 Balciauskas et al. 1999 Ulevicius et al. 1999 Ulevicius 2000 A. Ulevicius, pers. comm. |
| Luxembourg | 18th century? | – | 2000 | 1 ^s | Schley et al. 2001 |
| Mongolia/China | remnant | – | 1959-1985 | 800 | Lavrov & Hao-Tsuan 1961 Lavrov 1983 Stubbe & Dawaa 1983, 1986 |
| Netherlands | 1826 | – | 1988-2000 | 177-227 | Nolet 1994, 1996 J. Reinhold, pers. comm. H. Sluiter, pers. comm. |
| Norway | remnant | 1845 | 1925-1932 1952-1965 | c.70,000 | Djoshkin & Safonov 1972 Bevanger 1995 Rosell & Pedersen 1999 Andersen 2002 H. Parker, pers. comm. |
| Poland | 1844 | 1923 | 1943-1949 1975-1986 | 18,000- 23,000 | Zurowski & Kasperczyk 1986, 1988 Zurowski 1992 Macdonald et al. 1995 Czech 1999 Dzieciolowski & Gozdziwski 1999 A. Czech, pers. comm. |
| Romania | 1824? | – | 1998-1999 | >170 | Troidl & Ionescu 1997 G. Schwab, pers. comm. |

Table 2 (continued). The history and present status of European beavers (*Castor fiber*).

| Country | Extinction | Protection | Reintroduction and/or translocations | Present population size | References |
|-------------|--------------|------------|--------------------------------------|-------------------------|--|
| Russia | remnant | 1922 | 1927-1933 1934-1941 1946-1964 | 232,000- 300,000 | Djoshkin & Safonov 1972 Lavrov 1983 Dezhkin 1999 Safonov & Saveljev 1999 Saveljev & Safonov 1999 |
| Scotland | 16th century | – | ? | 0 ⁶ | Kitchener & Conroy 1997 Scottish Natural Heritage 2000 |
| Serbia | 1903? | – | 2004 | 30 | G. Schwab, pers. comm. |
| Slovenia | ? | ? | 1999 | <6 ⁷ | M. Grubestic, pers. comm. |
| Slovakia | 1851 | – | 1995 | >500 | Dúha & Majzlan 1997 Valachovic 1997 Pachinger & Hulik 1999 |
| Spain | 17th century | 1981 | 2003 | 18 | H. Lea, pers. comm. |
| Sweden | 1871 | 1873 | 1922-1939 | >100,000 | Freye 1978 Hartman 1994, 1995 |
| Switzerland | 1820 | – | 1956-1977 | >350 | Stocker 1985 Macdonald et al. 1995 Winter 1997 S. Capt, pers. comm. |
| Ukraine | remnant | 1922 | – | 6000 | Djoshkin & Safonov 1972 Lavrov & Lavrov 1986 Safonov & Saveljev 1999 |
| Wales | 12th century | – | – | 0 | Macdonald et al. 1995 |

¹ Spread into Bosnia-Herzegovina along the Sava river (from the Croatian reintroduction) is likely to have occurred, but no data.

² Based on subfossil remains. Philological evidence from placenames suggests a remnant may have survived as late as the 11th century.

³ Five animals reintroduced to a fenced enclosure at Ham Fen, Kent, 2003.

⁴ Finland also has a population of 14,000 American beavers (*Castor canadensis*).

⁵ Natural spread from Belgium.

⁶ Scottish Natural Heritage has applied to the Scottish Executive for permission to conduct a trial reintroduction in Knapdale, west Scotland. A decision is pending.

⁷ Natural spread from Croatia.

However, the longest distance *colonisation* to have been reported is apparently Hartman's (1995) record of colonisation of an area about 70 km away from the nearest other occupied area. Some time later, depending on the size and topography of the watershed but often after about 10-25 years, populations reach a critical density for encountering a mate, and the population then increases very rapidly. This is followed (on average 25-34 years after watershed colonisation in Hartman's (1995; personal communication) study) by a phase of population decline as marginal habitats become exhausted; and then by rough stability.

While in established populations average dispersal distance is usually much less than the extremes mentioned above, 3.9 km on the Azas river in Siberia, individuals disperse much further, one subadult male being found 85 km upstream (Saveljev et al. 2002). This implies the potential for considerable gene flow within continuous populations.

Incidences of beaver conflicts with humans tend to intensify during the later stages of the rapid increase phase, in part because then beaver more often take into use more marginal sites requiring more beaver engineering, dams, canals, etc, which may conflict with human landuses. In many countries, this phase of population development has been accompanied by the introduction of hunting, aimed at least in part at addressing conflict issues (Halley & Rosell 2002).

Conversely, while beavers can and do cross land, and have been found up to 11.7 km away from the nearest water body (Saveljev et al. 2002) watershed divisions do show a clear barrier effect for beaver expansion, which can be strongly isolating where natural or artificial habitat barriers, such as high mountains or intensive farmland, intrude between watersheds. Depending on the management strategy, therefore, this suggests a policy of many reintroductions to many watersheds, or, conversely, the early removal of colonising individuals on watersheds where their presence is considered undesirable. Given the pattern of range expansion within watersheds,

confining beaver populations to a particular stretch within a watershed will be impractical unless there are strong artificial barriers to expansion, such as man-made river barrages, or a heavy and directed hunting or trapping effort (Halley & Rosell 2002). Beaver populations should therefore be managed on a watershed scale.

Acknowledgements: Many beaver researchers throughout Europe kindly responded to requests for information. In particular, we would like to thank S. Asbirk, O. Boszér, K. Bevanger, S. Capt, D. Cirovic, A. Czech, R. Dennis, J.-F. DuCroz, M. Grubestic, G. Hartman, V. Kostkan, P. Lahti, H. Lea, R. Liobis, P. Rouland, A. Saveljev, G. Schwab, J. Sieber, F. Tattersall, U. Timm, A. Ulevicius and C. Winter.

Afterword: The authors would be grateful, on a continuing basis, for any corrections and/or updates to the information we have regarding the progress of reintroduction, range expansion, and population development of European beavers. All contributions will, of course, be acknowledged in any resulting publications.

References

- Andersen, O. 2002. Management of small game in Oppland, hunting season 2000/2001. NINA Research Report 734. Norwegian Institute for Nature Research, Trondheim, Norway. (In Norwegian)
- Balčiauskas, L., G. Trakimas, R. Juskaitis, A. Ulevicius & L. Balčiauskiene 1999. Atlas of Lithuanian mammals, amphibians and reptiles. Akstis Publications, Vilnius, Lithuania. (In Lithuanian with English summary)
- Balodis, M. 1992. Die Biber in Lettland. In: R. Schröpfer, M. Stubbe & D. Heidecke (eds.). Materialien des 2. Internationalen Symposiums Semiaquatische Säugetiere: 121-129. Martin-Luther-Universität, Halle/Saale, Germany.
- Balodis, M. 1994. Beaver population of Latvia: history, development and management. Proceedings of the Latvian Academy of Sciences, Section B 7-8: 122-127.
- Balodis, M. 1995. Beavers in Latvia. In: A. Ermala & S. Lahti (eds.). Proceedings of the Third Nordic Beaver Symposium, 1992, Finland: 6-9. Finnish Game and Fisheries Research Institute, Helsinki, Finland.
- Balodis, M. 1997. Recovery of the beaver in the in-

- constant Latvian landscape. In: K. Pachinger (ed.). Proceedings European Beaver Symposium, 15-19 September 1997: 3. Comenius University, Bratislava, Slovakia.
- Balodis, M. 1998. Beavers in the Gauja national park, Latvia. Proceedings of the Latvian Academy of Sciences 52: 49-53.
- Berthelsen, J.P. & A.B. Madsen 2003. Overvågning af bæver *Castor fiber* på Klosterheden Statsskovsdistrikt og vandløbssystemer i oplandet 2002. Arbejdsrapport no. 186, 2002. Danmarks miljøundersøgelser, Copenhagen, Denmark.
- Bevanger, K. 1995. Beverens gjenerobring av Norge. In: K. Brox (ed.). Natur 1995: 1-16. Tapir Forlag, Trondheim, Norway.
- Caughley, G. & A.R.E Sinclair 1994. Wildlife ecology and management. Blackwell Science, Cambridge, UK.
- Campbell, R. & F. Tattersall 2003. The Ham Fen beaver project – First Report. Wildlife Conservation Research Unit, University of Oxford, Oxford, UK.
- Collett, R. 1897. Bæveren i Norge, dens Utbredelsen og Levemaade 1896. Bergens Museums Aarbog 1: 1-139.
- Czech, A. 1999. The status of the European beaver in Poland. In: Proceedings of the Third International Symposium Semi-aquatic mammals and their habitats, 25-27 May 1999. Universität Osnabrück, Osnabrück, Germany.
- Danilov, P.I. 1995. Canadian and European beavers in the Russian northwest. In: A. Ermala & S. Lahti (eds.). Proceedings of the Third Nordic Beaver Symposium, 1992, Finland: 10-16. Finnish Game and Fisheries Research Institute, Helsinki, Finland.
- Dezhkin, V. 1999. Beaver in the modern world: necessity of the population management on the national and the international level. In: Proceedings of the First European-American Beaver Congress, Volga-Kama National Nature Preserve, 24-28 August 1999: 43-47. Special Publication of the Volga-Kama National Nature Preserve, Tatarstan, Russia.
- Djoshkin, W.W. & W.G. Safonov 1972. Die Biber der alten und neuen Welt. A. Ziemsen Verlag, Wittenberg, Germany.
- Dúha, J. & O. Majzlan 1997. The first reintroduction of beaver in Horná Orava in Slovakia. In: K. Pachinger (ed.). Proceedings European Beaver Symposium, 15-19 September 1997: 7. Comenius University, Bratislava, Slovakia.
- Dzieciolowski, R. & J. Gozdziwski 1999. The reintroduction of the European beaver *Castor fiber* in Poland: a success story. In: P. Busher & R. Dzieciolowski (eds.). Beaver protection, management and utilization in Europe and North America: 31-35. Kluwer Academic/Plenum Publishers, New York, USA.
- Ellegren, H., G. Hartman, M. Johansson & L. Andersson 1993. Major histocompatibility complex monomorphism and low levels of DNA fingerprinting variability in a reintroduced and rapidly expanding population of beavers. Proceedings National Academy Science USA 90: 8150-8153.
- Ermala, A, S. Lahti & P. Vikberg 1999. Bäverstammen ökar fortfarande – fangsten redan närmare 2500 bävvar. Jägaren 4: 28-31.
- Freye, H.A. 1978. *Castor fiber* Linnaeus, 1758 – Europäische Biber. In: J. Niethammer & F. Krapp (eds.). Handbuch der Säugetiere Europas: 184-200. Akademische Verlagsgesellschaft, Wiesbaden, Germany.
- Fustec, J., T. Lode, D. Le Jacques & J.P. Cormier 2001. Colonization, riparian habitat selection, and home range size in a reintroduced population of European beavers in the Loire. Freshwater Biology 46: 1361-1371.
- Halley, D.J. & F. Rosell 2002. The beaver's reconquest of Eurasia: Status, population development, and management of a conservation success. Mammal Review 32: 153-178.
- Hartman, G. 1994. Long-term population development of a reintroduced population in Sweden. Conservation Biology 8: 713-717.
- Hartman, G. 1995. Patterns of spread of a reintroduced beaver *Castor fiber* population in Sweden. Wildlife Biology 1: 97-103.
- Hartman, G. 1999. Beaver management and utilization in Scandinavia. In: P. Busher & R. Dzieciolowski (eds.). Beaver protection, management and utilization in Europe and North America: 1-7. Kluwer Academic/Plenum Publishers, New York, USA.
- Heideke, D. & K.-J. Horig 1986. Bestands- und Schutzsituation des Elbebibers. Halle & Magdeburg 23: 1-14.
- Heidecke, D. & P. Ibe 1997. Der Elbebiber. Biologie und Lebensweise. Biosphärenreservat "Mittlere Elbe", Dessau, Germany.
- Kitchener, A.C. & J.W.H. Conroy 1997. The history of the Eurasian beaver in Scotland. Mammal Review 27: 95-108.
- Kollar, H.P. & M. Seiter 1990. Biber in den Donau-Auen östlich von Wien. Eine erfolgreiche Wiedersiedlung. Verein für Ökologie und Umweltforschung, Wien, Austria.
- Kostkan, V. 1999. The European beaver, *Castor fiber* L. population growth in the Czech Republic. In: Proceedings of the Third International Symposium

- um Semi-aquatic mammals and their habitats, 25-27 May 1999. Universität Osnabrück, Osnabrück, Germany.
- Kostkan, V. & J. Lehky 1997. The Litovleské Pomoraví floodplain forest as a habitat for the reintroduction of the European beaver *Castor fiber* into the Czech Republic. *Global Ecology and Biogeography Letters* 6: 307-310.
- Laanetu, N. 1995. The status of European beaver *Castor fiber* L. population in Estonia and its influence on habitats. In: A. Ermala & S. Lahti (eds.). *Proceedings of the Third Nordic Beaver Symposium, 1992, Finland*: 34-40. Finnish Game and Fisheries Research Institute, Helsinki, Finland.
- Lavrov, L.S. 1980. Beaver. In: *Results of tagging animals*: 140-146. Nauka, Moscow, Russia. (In Russian)
- Lavrov, L.S. 1983. Evolutionary development of the genus *Castor* and taxonomy of the contemporary beavers of Eurasia. *Acta Zoologica Fennica* 174: 87-90.
- Lavrov, L.S. & L. Hao-Tsuan 1961. Present conditions and ecological peculiarities of beaver in natural colonies in Asia. *Vestnik Leningradskogo Universiteta* 9: 72-83.
- Lavrov, L.S. & V.L. Lavrov 1986. Verteilung und Anzahl ursprünglicher und aborigener Biberpopulationen und den USSR. *Zoologische Abhandlungen Dresden* 41: 105-109.
- Macdonald, D.W., F.H. Tattersall, E.D. Brown & D. Balharry 1995. Reintroducing the European beaver to Britain: nostalgic meddling or restoring biodiversity? *Mammal Review* 25: 161-200.
- Mickus, A. 1995. The European beaver *Castor fiber* L. in Lithuania. In: A. Ermala & S. Lahti (eds.). *Proceedings of the Third Nordic Beaver Symposium, 1992, Finland*: 44-45. Finnish Game and Fisheries Research Institute, Helsinki, Finland.
- Niewold, F.J.J. & D.D. Lammertsma 2000. Ruim 10 jaar bevers in de Biesbosch – een evaluatie van de populatieontwikkeling tot 2000. Report 15. Alterra, Wageningen, The Netherlands.
- Nolet, B.A. 1994. Return of the beaver to the Netherlands: viability and prospects of a re-introduced population. PhD-thesis. Rijksuniversiteit Groningen, Groningen, The Netherlands.
- Nolet, B.A. 1996. Management of the beaver *Castor fiber*: towards restoration of its former distribution and ecological function in Europe. Council of Europe/Convention on the Conservation of European Wildlife and Natural Habitats Bern Convention Nature & Environment series 86. Council of Europe Publishing, Strasbourg, France.
- Nolet, B. & F. Rosell 1998. Comeback of the beaver *Castor fiber*: an overview of old and new conservation problems. *Biological Conservation* 83: 165-173.
- Office Nationale de la Chasse 1997. Le castor dans le sud-est de la France. Office Nationale de la Chasse, Paris, France.
- Ozolins, J & J. Baumanis 2000. The current beaver status in Latvia. In: A. Czech & G. Schwab (eds.). *The European beaver in a new millenium. Proceedings of the 2nd European Beaver Symposium, 27-30 September 2000, Bialowieza*: 29. Carpathian Heritage Society, Krakow, Poland.
- Pachinger, K, & T. Hulik 1999. Origin, present conditions, and future prospects of the Slovakian beaver population. In: P. Busher & R. Dzieciolowski (eds.). *Beaver protection, management and utilisation in Europe and North America*: 43-52. Kluwer Academic/Plenum Publishers, New York, USA.
- Palionene, A. 1965. The beaver in the Lithuanian S.S.R. *Acta Theriologica* 10: 111-116.
- Richard, P.B. 1985. Peculiarities of the ecology and management of the Rhodanian beaver *Castor fiber* L. *Zeitschrift für angewandte Zoologie* 72: 143-152.
- Richard, P.B. 1986. The status of the beaver in France. *Zoologische Abhandlungen Dresden* 41: 121-130.
- Rosell, F. & K.V. Pedersen 1999. Bever. *Landbruksforlaget*, Oslo, Norway.
- Safonov, V. & A. Saveljev 1999. Beavers in CIS: resources, translocation and trapping. In: P. Busher & Y. Gorshkov (eds.). *Proceedings of the First European-American Beaver Congress, 24-28 August 1999*: 22-39. Volga-Kama National Nature Preserve, Kazan, Russia.
- Saveljev, A.P. & V.G. Safonov 1999. The beaver in Russia and adjoining countries. Recent trends in resource changes and management problems. In: P. Busher & R. Dzieciolowski (eds.). *Beaver protection, management and utilization in Europe and North America*: 17-24. Kluwer Academic/Plenum Publishers, New York, USA.
- Saveljev, A.P., M. Stubbe, A. Stubbe, V.V. Unzhakov & S.V. Kononov 2002. Natural movements of tagged beavers in Tyva. *Russian Journal of Ecology* 33: 434-439.
- Schley, L, L. Schmitz & C. Schank 2001. First record of the beaver *Castor fiber* in Luxembourg since at least the 19th century. *Lutra* 44: 41-42.
- Schwab, G., W. Dietzen & G. Von Lossow 1994. Biber in Bayern: Entwicklung eines Gesamtkonzeptes zum Schutz des Bibers. *Schriftenreihe Bayerische Ladesamt für Umweltschutz* 128: 9-31.

- Scottish Natural Heritage 2000. Reintroducing the European beaver to Scotland. Press release, 17/3/00.
- Sieber, J. 1999. "City beavers" in downtown Vienna. In: Proceedings of the Third International Symposium Semi-aquatic mammals and their habitats, 25-27 May 1999. Universität Osnabrück, Osnabrück, Germany.
- Skov- og Naturstyrelsen Denmark 2000. Det første bæverunge i 1000 år. Press release, 6/8/00.
- Stocker, G. 1985. The beaver in Switzerland: biological and ecological problems of re-establishment. Report 242. Swiss Federal Institute of Forestry Research, Berne, Switzerland.
- Stubbe, M. & N. Dawaa 1983. Akklimatisering des Zentralasiatischen Bibers – *Castor fiber birulai* Senbrennikov 1929 – in der Westmongolei. Erforschung biologischer ressourcen der Mongolischen Volksrepublik 2: 3-92.
- Stubbe, M. & N. Dawaa 1986. Die autochthone zentralasiatische Biberpopulation. Zoologische Abhandlungen Dresden 41: 91-103.
- Troidl, C. & G. Ionescu 1997. Beaver project Romania – a reintroduction with special focus on anthropic factors. In: K. Pachinger (ed.). Proceedings European Beaver Symposium, 15-19 September 1997: 29. Comenius University, Bratislava, Slovakia.
- Ulevicius, A. 2000. Temporal changes in an abundant beaver *Castor fiber* population. In: A. Czech & G. Schwab (eds.). The European beaver in a new millennium. Proceedings of the 2nd European Beaver Symposium, 27-30 September 2000, Bialowieza: 55. Carpathian Heritage Society, Krakow, Poland.
- Ulevicius, A., A. Mickus & R. Maseikyte 1999. History and present status of semiaquatic mammals in Lithuania. In: Proceedings of the Third International Symposium Semi-aquatic mammals and their habitats, 25-27 May 1999. Universität Osnabrück, Osnabrück, Germany.
- Valachovic, D. 1997. Distribution of the beaver in Záhorie region west Slovakia. In: K. Pachinger (ed.). Proceedings European Beaver Symposium, 15-19 September 1997: 31. Comenius University, Bratislava, Slovakia.
- Winter, C. 1997. Reintroduction of the beaver in Switzerland – a temporary or lasting success? In: K. Pachinger (ed.). Proceedings European Beaver Symposium, 15-19 September 1997: 33. Comenius University, Bratislava, Slovakia.
- Zharkov, I.V. & V.E. Solokov 1967. The European beaver in the Soviet Union. Acta Theriologica 12: 27-46.
- Zurowski, W. 1992. European beaver reintroduction into lowland and mountain tributaries of the Vistula river. In: B. Bobek, K. Perzanowski & W.L. Regelin (eds.). Global trends in wildlife management transactions of the 16th Congress of the International Union of Wildlife Biologists: 163-166. Swiat Press, Krakow, Poland.
- Zurowski, W. & B. Kasperczyk 1986. Characteristics of the European beaver population in the Suwalski lakeland. Acta Theriologica 31: 311-325.
- Zurowski, W. & B. Kasperczyk 1988. Effects of reintroduction of European beaver in the lowlands of the Vistula basin. Acta Theriologica 33: 325-338.

Samenvatting

Populatie en verspreiding van Europese bevers (*Castor fiber*)

Na een afname in het begin van de twintigste eeuw tot ongeveer 1200 dieren in acht verschillende populaties, hebben de Europese bevers (*Castor fiber*) zich krachtig hersteld, zowel in verspreiding als populatiegrootte, door vermindering van de jacht, natuurlijke verspreiding en op veel plaatsen uitgevoerde herintroducties. Populaties hebben zich nu (2003) gevestigd in alle landen binnen hun voormalige natuurlijke verspreidingsgebied met uitzondering van Groot-Britannië, Portugal, Italië en de zuidelijke Balkanlanden (Griekenland, Albanië, Bulgarije, Macedonië; de status in Bosnië-Herzegovina is onzeker). In Azië zijn er aanzienlijke populaties in centraal Siberië, Kamchatka en de Amur en kleine relict-populaties elders in Siberië en in Xinjiang (China) / west Mongolië. De huidige minimum schatting van de populatieomvang is 639.000. Zowel de populaties als het verspreidingsgebied breiden zich snel uit. We presenteren kaarten die beknopt de huidige wereldverspreiding van de Europese bever weergeven, en de Europees-Aziatische verspreiding van de uitgezette Amerikaanse bever (*Castor canadensis*). Verder laten we tabellen zien met de meest recente populatieschattingen voor elk land.

Received: 28 October 2003

Accepted: 31 March 2004